

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number
WO 2004/048365 A1

(51) International Patent Classification⁷: **C07D 401/12**,
407/14, 403/12, 407/12, 239/48, 417/12, 403/04, 403/14,
401/14, A61K 31/506, A61P 35/00

(74) Agent: SUH, Young, J.; Chiron Corporation, Intellectual
Property Dept., P.O. Box 8097, Emeryville, CA 94662-
8097 (US).

(21) International Application Number:
PCT/US2003/037294

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
21 November 2003 (21.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/428,473 21 November 2002 (21.11.2002) US
60/438,568 7 January 2003 (07.01.2003) US
60/523,081 19 November 2003 (19.11.2003) US

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): CHI-
RON CORPORATION [US/US]; 4560 Horton Street,
Emeryville, CA 94608-2916 (US).

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): NUSS, John, M.
[US/US]; 4560 Horton Street, Emeryville, CA 94608-2916
(US). PECCHI, Sabina [IT/US]; 4560 Horton Street,
Emeryville, CA 94608-2916 (US). RENHOWE, Paul, A.
[US/US]; 4560 Horton Street, Emeryville, CA 94608-2916
(US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: 2,4,6-TRISUBSTITUTED PYRIMIDINES AS PHOSPHOTIDYLINOSITOL (PI) 3-KINASE INHIBITORS AND
THEIR USE IN THE TREATMENT OF CANCER



(I)

(57) Abstract: Compounds having formula I are provided where the variables have the values de-
scribed herein. (I). Pharmaceutical formulations include the compounds or pharmaceutically accept-
able salts thereof and a pharmaceutically acceptable carrier and combinations with other agents. A
method of treating a patient comprises administering a pharmaceutical formulation according to the
invention to a patient in need thereof.

WO 2004/048365 A1

2,4,6-TRISUBSTITUTED PYRIMIDINES AS PHOSPHOTIDYLINOSITOL (PI) 3-KINASE
INHIBITORS AND THEIR USE IN THE TREATMENT OF CANCER

FIELD OF THE INVENTION

This invention pertains generally to the treatment of diseases, such as cancer,
5 characterized by the abnormal activity of growth factors, protein serine/threonine kinases,
and phospholipid kinases. In other aspects, the present invention provides small molecule
inhibitors of phosphatidylinositol (PI) 3-kinase, pharmaceutical formulations containing
such inhibitors, methods of treating patients with such pharmaceutical formulations, and
to methods of preparing such pharmaceutical formulations and inhibitors.

10 BACKGROUND OF THE INVENTION

Phosphatidylinositol 3-kinase (PI3K) is both a phospholipid kinase, and a protein
serine/threonine kinase as described in Carpenter *et al*, *Mol. Cell. Biol.* 13:1657-1665
(1993). PI3K is an enzyme stimulated by growth factors that is responsible for
phosphorylating phosphatidylinositol (PI) at the D-3' position of the inositol ring as
15 described in Whitman *et al*, *Nature* 332:644-646 (1988). PI3K association with Src-like
or receptor tyrosine kinases also implicates PI3K in the oncogenic or mitogenic responses
induced by these protein kinases, as described in Cantley *et al*, *Cell* 64:281-302 (1991),
Escobedo and Williams, *Nature* 335:85-87 (1988), and Fantl *et al*, *Cell* 69:413-423
(1992).

20 Previously, studies to elucidate the downstream effects of PI 3-kinase activation
have been conducted with receptor mutants constructed to alter the signal transduction of
PI3K, or by constructing mutant oncogenes to study a PI3K inducible oncogenic
response. The failure of receptor mutants of platelet derived growth factor (PDGF)
receptor to activate PI3K has been correlated with deficiency of the receptor mutants in
25 triggering a mitogenic response. Similarly, mutants of certain oncogenes have failed to
trigger the oncogenic transformation inducible by the parent oncogene. A method was
subsequently constructed to facilitate downstream effects of PI3K directly, without
growth factor activation to determine whether PI3K was distinctly involved oncogenesis
and mitogenesis. The results elucidated that PI3K can be directly or indirectly
30 responsible for many cellular processes, such as mitogenesis and oncogenesis, as well as
histamine secretion, neutrophil activation, platelet activation, cell migration, glucose
transport, antilipolysis, and vesicle sorting.


With the many regulatory responses associated with PI3-kinase, which is known to be involved in signal cascades involving other well known oncogenic proteins, such as receptor tyrosine kinases (e.g., VEGF-RTK), it would be highly desirable to produce small molecules capable of modulating, e.g. inhibiting, the activity of PI3-kinase.

It is an object of this invention to provide potent inhibitors of PI3K. It is further an object of the instant invention to provide compounds alone or in combination with other known agents to modulate cellular proliferation in patients in need thereof. Additionally, it is an object of this invention to provide medicaments for use in the treatment cancer.

SUMMARY OF THE INVENTION

The present invention provides novel pyrimidine based compounds, pharmaceutical formulations comprising the compounds, methods of inhibiting phosphatidylinositol 3-kinase (PI3K), and methods of treating cancer.

In one aspect, the present invention provides compounds of formula (I):




its stereoisomers, tautomers, pharmaceutically acceptable salts, esters, and prodrugs, wherein

Y is selected from the groups consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,

- (4) -O-,
 (5) -S-,
 (6) -SO-,
 (7) -SO₂-,
 (8) -C(R^{2x}, R^{3x})-, and
 (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
 (b) substituted or unsubstituted C₁-C₆-alkyl,
 (c) substituted or unsubstituted C₂-C₆-alkenyl,
 (d) substituted or unsubstituted C₂-C₆-alkynyl,
 (e) substituted or unsubstituted aryl,
 (f) substituted or unsubstituted heterocyclyl,
 (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
 (2) substituted or unsubstituted C₁-C₆-alkyl,
 (3) -COOH,
 (4) halo,
 (5) -OR^{1t}, and
 (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
 (2) substituted or unsubstituted heteroaryl, and
 (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
 (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^z is H or substituted or unsubstituted alkyl group; and

R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C_1 - C_6 -alkyl; and r is 0, 1, or 2;

with the proviso that when X is O, then Y is substituted or unsubstituted aryl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heteroaryl;

with the proviso that when W is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido morpholino, piperazino, or N-substituted piperazino, R₂ is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido-thiomorpholino, piperazino, or N'-[acetyl(alkanoyl of 1 to 3 carbon atoms)]piperazino, and X is NH, then

Y is not hydrogen, alkyl of 1 to 3 carbon atoms, cyclohexyl, phenyl, chloro-phenyl, carboxy-phenyl, carbomethoxy-phenyl, or pyridyl;

with the proviso that when W is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido morpholino, piperazino, or N-substituted piperazino, R₂ is
 5 morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido-thiomorpholino, piperazino, or N'-[acetyl(alkanoyl of 1 to 3 carbon atoms)]piperazino, and X is a direct link, then Y is not phenyl, substituted or unsubstituted C₁-C₆-alkyl, or 1-oxidothiomorpholino; and


with the proviso that when R₂ is phenyl independently substituted with one to five
 10 substituents selected from hydrogen, cycloalkyl, heterocycloalkyl, halo, nitro, amino, sulphonamido, or alkylsulphonylamino, R₁ is hydrogen, haloalkyl, alkyl, or halo, and X is NR^{1x}, then Y is substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocyclyl.

In one embodiment, the invention provides compounds of formula (I), wherein
 15 Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted aryl,
- (3) substituted or unsubstituted heterocyclyl, and
- (4) substituted or unsubstituted heteroaryl;


20 X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-, and



25 wherein R^{1x}, R^{2x}, R^{3x} are independently H or substituted or unsubstituted C₁-C₆-alkyl; and

W is selected from the group consisting of




wherein Z is -O- or -NR^z-, wherein R^{4w} is H or substituted or unsubstituted
 30 C₁-C₆-alkyl.

In another embodiment, the invention provides compounds of formula (I), wherein

Y is selected from the groups consisting of


- (1) substituted or unsubstituted heterocyclyl,
- (2) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and
- (4) ,

wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl; and

W is selected from the group consisting of




- wherein Z is $-O-$ or $-NR^Z-$, wherein R^{4w} is H or substituted or unsubstituted C_1 - C_6 -alkyl.

In another embodiment, the invention provides compounds of formula (I), wherein


Y is substituted or unsubstituted aryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and
- (4) ,

wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl; and

W is selected from the group consisting of



wherein Z is -O- or -NR^Z, wherein R^{4w} is H or substituted or unsubstituted C₁-C₆-alkyl.

In another embodiment, the invention provides compounds of formula (I), wherein


5 Y is substituted or unsubstituted alkyl;

X is selected from the group consisting of

(1) a direct link,


(2) -N(R^{1x})-,

(3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-, and

10 (4) ,

wherein R^{1x}, R^{2x}, R^{3x} are independently H or substituted or unsubstituted C₁-C₆-alkyl; and

W is selected from the group consisting of



15 wherein Z is -O- or -NR^Z, wherein R^{4w} is H or substituted or unsubstituted C₁-C₆-alkyl.

In another embodiment, the invention provides compounds of formula (I), wherein

Y is selected from the group consisting of

20 (1) substituted or unsubstituted heterocyclyl,


(2) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

(1) a direct link,

(2) -N(R^{1x})-,


25 (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-, and

(4) ,

wherein R^{1x}, R^{2x}, R^{3x} are independently H or substituted or unsubstituted C₁-C₆-alkyl;

R₂ is substituted or unsubstituted aryl; and

W is



wherein Z is -O- or -NH-.


In another embodiment, the invention provides compounds of formula (I),

5 wherein

Y is substituted or unsubstituted aryl;

X is selected from the group consisting of


- (1) a direct link,
- (2) -N(R^{1x})-,
- 10 (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-, and



wherein R^{1x}, R^{2x}, R^{3x} are independently H or substituted or unsubstituted C₁-C₆-alkyl;

R₂ is substituted or unsubstituted aryl; and

15 W is



wherein Z is -O- or -NH-.


In another embodiment, the invention provides compounds of formula (I),

wherein

20 Y is substituted or unsubstituted alkyl;

X is selected from the group consisting of


- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-, and



wherein R^{1x}, R^{2x}, R^{3x} are independently H or substituted or unsubstituted C₁-C₆-alkyl;


R₂ is substituted or unsubstituted aryl; and

W is



wherein Z is -O- or -NH-.


In another embodiment, the invention provides compounds of formula (I) having
5 structure (II):




wherein Y is selected from the group consisting of

- 10
- (1) substituted or unsubstituted C₁-C₆-alkyl,
 - (2) substituted or unsubstituted aryl,
 - (3) substituted or unsubstituted heterocyclyl, and
 - (4) substituted or unsubstituted heteroaryl; and


X is selected from the group consisting of

- 15
- (1) a direct link,
 - (2) -N(R^{1x})-,
 - (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-, and
 - (4) .


In another embodiment, the invention provides compounds of formula (I) having structure (II), wherein Y and X, taken together, are selected from the group consisting of




In another embodiment, the invention provides compounds of formula (I) having structure (II), wherein Y and X, taken together, are selected from the group consisting of



In another embodiment, the invention provides compounds of formula (I) having structure (II), wherein Y and X, taken together, are selected from the group consisting of



In another embodiment, the invention provides compounds of formula (I) having structure (III):




(III)

wherein R_3 , R_4 , R_5 , R_6 are selected from the group consisting of

- (1) H ,
- (2) substituted or unsubstituted C_1 - C_6 -alkyl,
- (3) $-\text{COOR}^{\text{lt}}$,
- (4) $-\text{COONH}_2$,
- (5) $-\text{OR}^{\text{lt}}$, and
- (6) $-\text{NHR}^{\text{lt}}$.

In another embodiment, the invention provides compounds of formula (I) having structure (IV):



(IV)

wherein R_3 , R_4 , R_5 , R_6 are selected from the group consisting of


- (1) H ,
- (2) substituted or unsubstituted C_1 - C_6 -alkyl,
- (3) $-\text{COOR}^{\text{lt}}$,
- (4) $-\text{COONH}_2$,

(5) -OR^{1t}, and

(6) -NHR^{1t}.

In another embodiment, the invention provides compounds of formula (I) having structure (V):

5



(V)

wherein R₃, R₄, R₅, R₆ are selected from the group consisting of

- 10
- (1) H,
 - (2) substituted or unsubstituted C₁-C₆-alkyl,
 - (3) -COOR^{1t},
 - (4) -COONH₂,
 - (5) -OR^{1t}, and
 - (6) -NHR^{1t}; and

15 R^{2a} and R^{2b} are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) halo,
- (4) -(CH₂)_q-N(R^{2c}, R^{2d}),
- 20 (5) -(CH₂)_q-N(R^{2c}, R^{2d})COR^{2e},
- (6) -(CH₂)_q-OR^{2e},
- (7) -(CH₂)_q-OCOR^{2e},
- (8) -(CH₂)_q-OCOOR^{2e},
- (9) -(CH₂)_q-COOR^{2e},
- 25 (10) -(CH₂)_q-CONR^{2c},
- (11) -CN,
- (12) -NO₂,
- (13) -SO₂NH₂,
- (14) -NHSO₂CH₃, and

(15) $-\text{SO}_2\text{R}^{2f}$,

wherein R^{2c} , R^{2d} , R^{2e} , and R^{2f} are selected from the group consisting of


(a) H,

(b) substituted or unsubstituted alkyl, and

(c) substituted or unsubstituted phenyl; and


q is 0, 1, 2, 3, or 4.

In another embodiment, the invention provides compounds of formula (I) having structure (VI):




(VI)

wherein R_2 is selected from the group consisting of



In another embodiment, the invention provides compounds of formula (I) having structure (VII):




(VII)

wherein R₇, R₈, R₉, and R₁₀ are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},
- (4) -COONH₂,
- (5) -OR^{1t}, and
- (6) -NHR^{1t}.

In another embodiment, the invention provides compounds of formula (I) having structure (VIII):




(VIII)

wherein R₇, R₈, R₉, R₁₀ are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},
- (4) -CONH₂,
- (5) -OR^{1t}, and
- (6) -NHR^{1t}.

In another embodiment, the invention provides compounds of formula (I) having structure (IX):



(IX)

wherein R^{1a} and R^{1b} are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) halo,
- (4) $-(CH_2)_q-N(R^{2c}, R^{2d})$,
- (5) $-(CH_2)_q-N(R^{2c}, R^{2d})COR^{2e}$,
- (6) $-(CH_2)_q-OR^{2e}$,
- (7) $-(CH_2)_q-OCOR^{2e}$,
- (8) $-(CH_2)_q-OCOOR^{2e}$,
- (9) $-(CH_2)_q-COOR^{2e}$,
- (10) $-(CH_2)_q-CONR^{2c}$,
- (11) -CN,
- (12) -NO₂,
- (13) -SO₂NH₂,
- (14) -NHSO₂CH₃, and
- (15) -SO₂R^{2f},

wherein R^{2c} , R^{2d} , R^{2e} , and R^{2f} are selected from the group consisting of


- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted phenyl; and

R_7 is selected from the group consisting of


- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},

- (4) $-\text{COONH}_2$,
- (5) $-\text{OR}^{\text{1t}}$, and
- (6) $-\text{NHR}^{\text{1t}}$.


In another embodiment, the invention provides compounds of formula (I) having
5 structure (X):



wherein R_2 is selected from the group consisting of





In another embodiment, the invention provides compounds of formula (I) having structure (XI):



(XI)

wherein R^{2g} is selected from the group consisting of


- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) -CONHR^{2h},
- (4) -CON(R^{2h})-(CH₂)₂₋₃-N(R^{2h}, R²ⁱ),
- (5) -COR^{2j},
- (6) -CO₂R^{2j},
- (7) -COC₁₋₆-alkyl-CO₂H,
- (8) -CH₂-OC(=O)R²ⁱ,
- (9) -CH₂-OC(=O)NHCHR²ⁱCO₂R^{2j},
- (10) -P(=O)(OR^{2k}, OR^{2p}),



wherein R^{2h}, R²ⁱ, R^{2j}, R^{2k}, and R^{2p} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted aryl.



In another embodiment, the invention provides compounds of formula (I) having structure (XII):



(XII)

wherein R^{2g} is selected from the group consisting of


- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) -CONHR^{2h},
- (4) -CON(R^{2h})-(CH₂)₂₋₃-N(R^{2h}, R²ⁱ),
- (5) -COR^{2j},
- (6) -CO₂R^{2j},
- (7) -COC₁₋₆-alkyl-CO₂H,
- (8) -CH₂-OC(=O)R²ⁱ,
- (9) -CH₂-OC(=O)NHCHR²ⁱCO₂R^{2j},
- (10) -P(=O)(OR^{2k}, OR^{2p}),




wherein R^{2h}, R²ⁱ, R^{2j}, R^{2k}, and R^{2p} are selected from the group consisting

- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted aryl.

In another embodiment, the present invention provides compounds of the following formula (I):



5 wherein W is:



wherein Z is selected from the group consisting of -CH₂-, -NH-, -O-, -S-, and -NR₆-, where R₆ is an alkyl or substituted alkyl group;

R₃ is absent or selected from the group consisting of alkyl, substituted alkyl, amino, alkylamino, aminoalkyl, dialkylamino, dialkylaminoalkyl, alkoxy, alkenyl, substituted alkenyl, alkynyl, carbonylamino, and alkoxycarbonyl; and

m and n are integers from 0-2;

X is a covalent bond or is selected from the group consisting of -CH₂-, -CHF-, -CF₂-, -NH-, -O-, -S-, and -NR₅-, where R₅ is an alkyl or substituted alkyl group;

15 Y is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;


R₁ is selected from the group consisting of hydrogen, halogen, carboxylic acid, and alkyl; and

20 R₂ is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

the tautomers thereof;

and the pharmaceutically acceptable salts, esters, or prodrugs thereof.

In another aspect of the compound of formula (I), W is a morpholinyl group as shown below:



25

wherein, R_3 , m , and n are as described above.

In another more particular embodiment of compound (I), W is an unsubstituted morpholinyl group.

In another more particular embodiment of compound (I), X is $-NH-$.

5 In another more particular embodiment of compound (I), Y is a heteroaryl or substituted heteroaryl group selected from pyridyl, and alkoxy pyridyl.


In another more particular embodiment of compound (I), R_1 is hydrogen.

In another more particular embodiment of compound (I), R_2 is an aryl or substituted aryl group.

10 In another more particular embodiment of compound (I), R_2 is selected from the group consisting of phenyl, phenol, aniline, hydroxybenzyl, phenylalkoxycarbonyl, phenylcarbonylalkoxy, phenylaminocarbonyl, and phenylcarbonylamino.

In another more particular embodiment of compound (I), R_3 is absent.

In another embodiment, compounds of formula (II) are provided:



wherein X is selected from the group consisting of $-NH-$, $-O-$, and $-S-$;

Y is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

20 R_1 is hydrogen, halogen, or a carboxylic acid group;

R_2 is selected from the group consisting of aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

the tautomers thereof;

and the pharmaceutically acceptable salts, esters, or prodrugs thereof.

25 In another aspect of the compound of formula (II), X is $-NH-$.

In another aspect of the compound of formula (II), Y is a heteroaryl or substituted heteroaryl group selected from pyridyl, and alkoxy pyridyl.


In another aspect of the compound of formula (II), R_1 is absent.

In another aspect of the compound of formula (II), R_2 is an aryl or substituted aryl group.

In another aspect of the compound of formula (II), R_2 is selected from the group consisting of phenyl, phenol, aniline, hydroxybenzyl, phenylalkoxycarbonyl, phenylcarbonylalkoxy, phenylaminocarbonyl, and phenylcarbonylamino.

In another aspect of the compound of formula (II), R_3 is absent.

In another embodiment, compounds of formula (XIII) are provided:



XIII

wherein,

X is selected from the group consisting of -NH-, -O-, and -S-;

Y is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

R_1 is hydrogen, halogen, or a carboxylic acid;

R_4 is independently selected from the group consisting of hydrogen, halogen, alkyl, substituted alkyl, hydroxy, alkoxy, amino, alkylamino, aminoalkyl, dialkylamino, dialkylaminoalkyl, aryl, heteroaryl, heterocyclyl, carbonylamino, and alkoxycarbonyl; and

q is an integer from 1-5;

the tautomers thereof;

and the pharmaceutically acceptable salts, esters, or prodrugs thereof.

In another aspect of the compound of formula (XIII), X is -NH- and R_1 is hydrogen.

In another aspect of the compound of formula (XIII), R_4 is selected from the group consisting of hydrogen, halogen, alkyl, substituted alkyl, hydroxy, alkoxy, amino, alkylamino, aminoalkyl, dialkylamino, dialkylaminoalkyl, carbonylamino, and alkoxycarbonyl.

In addition to the compounds described above with the noted provisos, the present invention also includes compounds defined as above, but that do not include the noted provisos. This second class of compounds of the invention does include the following provisos:

5 when X is O, then Y is substituted or unsubstituted aryl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heteroaryl;

 when W is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido morpholino, piperazino, or N-substituted piperazino, R₂ is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido-thiomorpholino, piperazino, or N'-[acetyl(alkanoyl
10 of 1 to 3 carbon atoms)]piperazino, and X is NH, then Y is not hydrogen, alkyl of 1 to 3 carbon atoms, cyclohexyl, phenyl, chloro-phenyl, carboxy-phenyl, carbomethoxy-phenyl, or pyridyl;

 when W is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido morpholino, piperazino, or N-substituted piperazino, R₂ is morpholino, thiomorpholino,
15 1-oxido-thiomorpholino, 1,1-dioxido-thiomorpholino, piperazino, or N'-[acetyl(alkanoyl of 1 to 3 carbon atoms)]piperazino, and X is a direct link, then Y is not substituted or unsubstituted C1-C6-alkyl, or 1-oxidothiomorpholino; and

 when R₂ is phenyl independently substituted with one to five substituents selected from hydrogen, cycloalkyl, heterocycloalkyl, halo, nitro, amino, sulphonamido, or
20 alkylsulphonylamino, R₁ is hydrogen, haloalkyl, alkyl, or halo, and X is NR^{1x}, then Y is substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocyclyl.

Thus, in one aspect, the present invention provides two class of compounds, each including provisos. The difference between the two classes of compounds being the difference in provisos.

25 In other aspects, the invention provides methods for using compounds that are inhibitors of phosphatidylinositol 3-kinase (PI3K).

 In another aspect of the invention, pharmaceutical formulations are provided that include one or more of the compounds described herein in combination with a pharmaceutically acceptable carrier.

30 Further objects, features, and advantages of the invention will be apparent from the following detailed description.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention provides novel compounds that act as inhibitors of serine/threonine kinases, phospholipid kinases, and, more particularly, as inhibitors of phosphatidylinositol 3-kinase (PI3K) function. The compounds provided herein can be formulated into pharmaceutical formulations that are useful in treating patients with a need for an inhibitor of PI3K, especially, in particular embodiments, to provide compositions and methods for reducing cellular proliferation and in the treatment of cancer.

The following abbreviations and definitions are used throughout this application:

<u>Abbreviation</u>	<u>Meaning</u>
PI3K	phosphatidylinositol 3-kinase
AcOH	acetic acid
ATP:	adenosine triphosphate
BOC	tert-butoxycarbonyl
CPT 11	irinotecan
DIBAL-H	diisobutylaluminum hydride
DCM	dichloromethane
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
DIEA	diisopropylethylamine
DMA:	N,N-Dimethylacetamide
DMF:	N,N-Dimethylformamide
DMSO	dimethyl sulfoxide
EDTA:	ethylene diamine tetraacetic acid
EtOAc:	ethyl acetate
EtOH:	ethanol
5-FU	5-fluorouracil
GCMS	Gas Chromatography / Mass Spectroscopy
HBTU:	O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate
HPLC	High Performance Liquid Chromatography
IC ₅₀ value:	the concentration of an inhibitor that causes

<u>Abbreviation</u>	<u>Meaning</u>
	a 50 % reduction in a measured activity.
LCMS	Liquid Chromatography / Mass Spectroscopy
MeOH:	methanol
NMP:	N-methylpyrrolidone
NMR	nuclear magnetic resonance
Rt	room temperature (25°C)
THF:	tetrahydrofuran
TLC	thin-layer chromatography

The phrase "alkyl" refers to alkyl groups that do not contain heteroatoms. Thus the phrase includes straight chain alkyl groups such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl and the like. The phrase also includes branched chain isomers of straight chain alkyl groups, including but not limited to, the following which are provided by way of example: $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)_3$, $-\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, and others. The phrase also includes cyclic alkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl and such rings substituted with straight and branched chain alkyl groups as defined above. Thus the phrase alkyl groups include primary alkyl groups, secondary alkyl groups, and tertiary alkyl groups. Preferred alkyl groups include straight and branched chain alkyl groups and cyclic alkyl groups having 1 to 12 carbon atoms.

The phrase "substituted alkyl" refers to an alkyl group as defined above in which one or more bonds to a carbon(s) or hydrogen(s) are replaced by a bond to non-hydrogen and non-carbon atoms such as, but not limited to, a halogen atom such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, and ester groups; a sulfur atom in groups such as thiol groups, alkyl and aryl sulfide groups,

sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as in trialkylsilyl groups, dialkylarylsilyl groups, alkyl diarylsilyl groups, and triarylsilyl groups; and other
5 heteroatoms in various other groups. Substituted alkyl groups also include groups in which one or more bonds to a carbon(s) or hydrogen(s) atom is replaced by a higher-order bond (e.g., a double- or triple-bond) to a heteroatom such as oxygen in oxo, carbonyl, carboxyl, and ester groups; nitrogen in groups such as imines, oximes, hydrazones, and nitriles. Substituted alkyl groups further include alkyl groups in which
10 one or more bonds to a carbon(s) or hydrogen(s) atoms is replaced by a bond to an aryl, heterocyclyl group, or cycloalkyl group. Preferred substituted alkyl groups include, among others, alkyl groups in which one or more bonds to a carbon or hydrogen atom is/are replaced by one or more bonds to fluorine atoms. Another preferred substituted alkyl group is the trifluoromethyl group and other alkyl groups that contain the
15 trifluoromethyl group. Other preferred substituted alkyl groups include those in which one or more bonds to a carbon or hydrogen atom is replaced by a bond to an oxygen atom such that the substituted alkyl group contains a hydroxyl, alkoxy, or aryloxy group. Still other preferred substituted alkyl groups include alkyl groups that have an amine, or a substituted or unsubstituted alkylamine, dialkylamine, arylamine, (alkyl)(aryl)amine, diarylamine, heterocyclylamine, diheterocyclylamine, (alkyl)(heterocyclyl)amine, or
20 (aryl)(heterocyclyl)amine group.

By halo is meant chloro, bromo, iodo, or fluoro or by halogen is meant chlorine, bromine, iodine or fluorine.

The phrase "alkenyl" refers to straight and branched chain and cyclic groups such
25 as those described with respect to alkyl groups as defined above, except that at least one double bond exists between two carbon atoms. Examples include, but are not limited to vinyl, $-\text{CH}=\text{C}(\text{H})(\text{CH}_3)$, $-\text{CH}=\text{C}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)=\text{C}(\text{H})_2$, $-\text{C}(\text{CH}_3)=\text{C}(\text{H})(\text{CH}_3)$, $-\text{C}(\text{CH}_2\text{CH}_3)=\text{CH}_2$, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, and hexadienyl among others. The phrase "substituted alkenyl" has the same
30 meaning with respect to alkenyl groups that substituted alkyl groups has with respect to unsubstituted alkyl groups. A substituted alkenyl group includes alkenyl groups in which a non-carbon or non-hydrogen atom is bonded to a carbon double bonded to another

carbon and those in which one of the non-carbon or non-hydrogen atoms is bonded to a carbon not involved in a double bond to another carbon.

The phrase "alkynyl" refers to straight and branched chain groups such as those described with respect to alkyl groups as defined above, except that at least one triple bond exists between two carbon atoms. Examples include, but are not limited to
5 $-C\equiv C(H)$, $-C\equiv C(CH_3)$, $-C\equiv C(CH_2CH_3)$, $-C(H_2)C\equiv C(H)$, $-C(H)_2C\equiv C(CH_3)$, and $-C(H)_2C\equiv C(CH_2CH_3)$ among others. The phrase "substituted alkynyl" has the same meaning with respect to alkynyl groups that substituted alkyl groups had with respect to unsubstituted alkyl groups. A substituted alkynyl group includes alkynyl groups in which
10 a non-carbon or non-hydrogen atom is bonded to a carbon triple bonded to another carbon and those in which a non-carbon or non-hydrogen atom is bonded to a carbon not involved in a triple bond to another carbon.

The phrase "heterocyclyl" refers to both aromatic and nonaromatic ring compounds including monocyclic, bicyclic, and polycyclic ring compounds such as, but
15 not limited to, quinuclidinyl, containing 3 or more ring members of which one or more is a heteroatom such as, but not limited to, N, O, and S. Although the phrase "unsubstituted heterocyclyl" includes condensed heterocyclic rings such as benzimidazolyl, it does not include heterocyclyl groups that have other groups such as alkyl or halo groups bonded to one of the ring members as compounds such as 2-methylbenzimidazolyl are substituted
20 heterocyclyl groups. Examples of heterocyclyl groups include, but are not limited to: unsaturated 3- to 8-membered rings containing 1 to 4 nitrogen atoms such as, but not limited to pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, dihydropyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl etc.), tetrazolyl, (e.g., 1H-tetrazolyl, 2H-tetrazolyl, etc.); saturated
25 3- to 8-membered rings containing 1 to 4 nitrogen atoms such as, but not limited to, pyrrolidinyl, imidazolidinyl, piperidinyl, piperazinyl; condensed unsaturated heterocyclic groups containing 1 to 4 nitrogen atoms such as, but not limited to, indolyl, isoindolyl, indolinyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl; unsaturated 3- to 8-membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen
30 atoms such as, but not limited to, oxazolyl, isoxazolyl, oxadiazolyl (e.g. 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxa-diazolyl, etc.); saturated 3- to 8-membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms such as, but not limited to, morpholinyl; unsaturated condensed heterocyclic groups containing 1 to 2

oxygen atoms and 1 to 3 nitrogen atoms, for example, benzoxazolyl, benzoxadiazolyl, benzoxazinyl (e.g. 2H-1,4-benzoxazinyl etc.); unsaturated 3- to 8-membered rings containing 1 to 3 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolyl, isothiazolyl, thiadiazolyl (e.g. 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.); saturated 3- to 8-membered rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolodinyll; saturated and unsaturated 3- to 8-membered rings containing 1 to 2 sulfur atoms such as, but not limited to, thienyl, dihydrodithienyl, dihydrodithionyl, tetrahydrothiophene, tetra-hydrothiopyran; unsaturated condensed heterocyclic rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, benzothiazolyl, benzothiadiazolyl, benzothiazinyl (e.g. 2H-1,4-benzothiazinyl, etc.), dihydrobenzothiazinyl (e.g. 2H-3,4-dihydrobenzothiazinyl, etc.), unsaturated 3- to 8-membered rings containing oxygen atoms such as, but not limited to furyl; unsaturated condensed heterocyclic rings containing 1 to 2 oxygen atoms such as benzodioxolyl (e.g. 1,3-benzodioxoyl, etc.); unsaturated 3- to 8-membered rings containing an oxygen atom and 1 to 2 sulfur atoms such as, but not limited to, dihydrooxathienyl; saturated 3- to 8-membered rings containing 1 to 2 oxygen atoms and 1 to 2 sulfur atoms such as 1,4-oxathiane; unsaturated condensed rings containing 1 to 2 sulfur atoms such as benzothienyl, benzodithienyl; and unsaturated condensed heterocyclic rings containing an oxygen atom and 1 to 2 oxygen atoms such as benzoxathienyl. Heterocyclyl group also include those described above in which one or more S atoms in the ring is double-bonded to one or two oxygen atoms (sulfoxides and sulfones). For example, heterocyclyl groups include tetrahydrothiophene, tetrahydrothiophene oxide, and tetrahydrothiophene 1,1-dioxide. Preferred heterocyclyl groups contain 5 or 6 ring members. More preferred heterocyclyl groups include morpholine, piperazine, piperidine, pyrrolidine, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, thiomorpholine, thiomorpholine in which the S atom of the thiomorpholine is bonded to one or more O atoms, pyrrole, homopiperazine, oxazolidin-2-one, pyrrolidin-2-one, oxazole, quinuclidine, thiazole, isoxazole, furan, and tetrahydrofuran.

The phrase "substituted heterocyclyl" refers to a heterocyclyl group as defined above in which one of the ring members is bonded to a non-hydrogen atom such as described above with respect to substituted alkyl groups and substituted aryl groups. Examples, include, but are not limited to, 2-methylbenzimidazolyl,

5-methylbenzimidazolyl, 5-chlorobenzthiazolyl, 1-methyl piperazinyl, and 2-chloropyridyl among others.

The phrase "aryl" refers to aryl groups that do not contain heteroatoms. Thus the phrase includes, but is not limited to, groups such as phenyl, biphenyl, anthracenyl, naphthenyl by way of example. Although the phrase "unsubstituted aryl" includes groups containing condensed rings such as naphthalene, it does not include aryl groups that have other groups such as alkyl or halo groups bonded to one of the ring members, as aryl groups such as tolyl are considered herein to be substituted aryl groups as described below. A preferred unsubstituted aryl group is phenyl. Unsubstituted aryl groups may be bonded to one or more carbon atom(s), oxygen atom(s), nitrogen atom(s), and/or sulfur atom(s) in the parent compound, however.

The phrase "substituted aryl group" has the same meaning with respect to unsubstituted aryl groups that substituted alkyl groups had with respect to unsubstituted alkyl groups. However, a substituted aryl group also includes aryl groups in which one of the aromatic carbons is bonded to one of the non-carbon or non-hydrogen atoms described above and also includes aryl groups in which one or more aromatic carbons of the aryl group is bonded to a substituted and/or unsubstituted alkyl, alkenyl, or alkynyl group as defined herein. This includes bonding arrangements in which two carbon atoms of an aryl group are bonded to two atoms of an alkyl, alkenyl, or alkynyl group to define a fused ring system (e.g. dihydronaphthyl or tetrahydronaphthyl). Thus, the phrase "substituted aryl" includes, but is not limited to tolyl, and hydroxyphenyl among others.

The term "heteroaryl", as used herein, refers to a cyclic or bicyclic aromatic radical having from five to ten ring atoms in each ring of which one atom of the cyclic or bicyclic ring is selected from S, O and N; zero, one or two ring atoms are additional heteroatoms independently selected from S, O and N; and the remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms, such as, for example, pyridyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, and naphthyridinyl, and the like.

The term "substituted heteroaryl" as used herein refers to a heteroaryl group as defined herein substituted by independent replacement of one, two or three of the hydrogen atoms thereon with Cl, Br, F, I, -OH, -CN, C₁-C₃-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkoxy substituted with aryl, haloalkyl, thioalkoxy, amino, alkylamino,

dialkylamino, mercapto, nitro, carboxaldehyde, carboxy, alkoxycarbonyl and carboxamide. In addition, any one substituent may be an aryl, heteroaryl, or heterocycloalkyl group.

The term "biaryl" refers to a group or substituent to which two aryl groups, which are not condensed to each other, are bound. Exemplary biaryl compounds include, for example, phenylbenzene, diphenyldiazene, 4-methylthio-1-phenylbenzene, phenoxybenzene, (2-phenylethynyl)benzene, diphenyl ketone, (4-phenylbuta-1,3-diynyl)benzene, phenyl-benzylamine, (phenylmethoxy)benzene, and the like. Preferred optionally substituted biaryl groups include: 2-(phenylamino)-N-[4-(2-phenylethynyl)phenyl]acetamide, 1,4-diphenylbenzene, N-[4-(2-phenylethynyl)phenyl]-2-[benzylamino]acetamide, 2-amino-N-[4-(2-phenylethynyl)phenyl]propanamide, 2-amino-N-[4-(2-phenylethynyl)phenyl]acetamide, 2-(cyclopropylamino)-N-[4-(2-phenylethynyl)phenyl]acetamide, 2-(ethylamino)-N-[4-(2-phenylethynyl)phenyl]acetamide, 2-[(2-methylpropyl)amino]-N-[4-(2-phenylethynyl)phenyl]acetamide, 5-phenyl-2H-benzo[d]1,3-dioxolene, 2-chloro-1-methoxy-4-phenylbenzene, 2-[(imidazolylmethyl)amino]-N-[4-(2-phenylethynyl)phenyl]acetamide, 4-phenyl-1-phenoxybenzene, N-(2-aminoethyl)[4-(2-phenylethynyl)phenyl]carboxamide, 2-[[4-fluorophenyl)methyl]amino}-N-[4-(2-phenylethynyl)phenyl]acetamide, 2-[[4-methylphenyl)methyl]amino}-N-[4-(2-phenylethynyl)phenyl]acetamide, 4-phenyl-1-(trifluoromethyl)benzene, 1-butyl-4-phenylbenzene, 2-(cyclohexylamino)-N-[4-(2-phenylethynyl)phenyl]acetamide, 2-(ethylmethylamino)-N-[4-(2-phenylethynyl)phenyl]acetamide, 2-(butylamino)-N-[4-(2-phenylethynyl)phenyl]acetamide, N-[4-(2-phenylethynyl)phenyl]-2-(4-pyridylamino)acetamide, N-[4-(2-phenylethynyl)phenyl]-2-(quinuclidin-3-ylamino)acetamide, N-[4-(2-phenylethynyl)phenyl]pyrrolidin-2-ylcarboxamide, 2-amino-3-methyl-N-[4-(2-phenylethynyl)phenyl]butanamide, 4-(4-phenylbuta-1,3-diynyl)phenylamine, 2-(dimethylamino)-N-[4-(4-phenylbuta-1,3-diynyl)phenyl]acetamide, 2-(ethylamino)-N-[4-(4-phenylbuta-1,3-diynyl)phenyl]acetamide, 4-ethyl-1-phenylbenzene, 1-[4-(2-phenylethynyl)phenyl]ethan-1-one, N-(1-carbamoyl-2-hydroxypropyl)[4-(4-phenylbuta-1,3-diynyl)phenyl]carboxamide, N-[4-(2-phenylethynyl)phenyl]propanamide, 4-methoxyphenyl phenyl ketone, phenyl-N-benzamide, (tert-butoxy)-N-[(4-phenylphenyl)methyl]carboxamide, 2-(3-phenylphenoxy)ethanehydroxamic acid, 3-phenylphenyl propanoate, 1-(4-ethoxyphenyl)-4-methoxybenzene, and [4-(2-phenylethynyl)phenyl]pyrrole.

The term "heteroarylaryl" refers to a biaryl group where one of the aryl groups is a heteroaryl group. Exemplary heteroarylaryl groups include, for example, 2-phenylpyridine, phenylpyrrole, 3-(2-phenylethynyl)pyridine, phenylpyrazole, 5-(2-phenylethynyl)-1,3-dihydropyrimidine-2,4-dione, 4-phenyl-1,2,3-thiadiazole, 5 2-(2-phenylethynyl)pyrazine, 2-phenylthiophene, phenylimidazole, 3-(2-piperazinylphenyl)furan, 3-(2,4-dichlorophenyl)-4-methylpyrrole, and the like. Preferred optionally substituted heteroarylaryl groups include: 5-(2-phenylethynyl)pyrimidine-2-ylamine, 1-methoxy-4-(2-thienyl)benzene, 1-methoxy-3-(2-thienyl)benzene, 5-methyl-2-phenylpyridine, 5-methyl-3- 10 phenylisoxazole, 2-[3-(trifluoromethyl)phenyl]furan, 3-fluoro-5-(2-furyl)-2-methoxy-1-prop-2-enylbenzene, (hydroxyimino)(5-phenyl(2-thienyl))methane, 5-[(4-methylpiperazinyl)methyl]-2-phenylthiophene, 2-(4-ethylphenyl)thiophene, 4-methylthio-1-(2-thienyl)benzene, 2-(3-nitrophenyl)thiophene, (tert-butoxy)-N-[(5-phenyl(3-pyridyl))methyl]carboxamide, hydroxy-N-[(5-phenyl(3-pyridyl))methyl]amide, 15 2-(phenylmethylthio)pyridine, and benzylimidazole.


The term "heteroarylheteroaryl" refers to a biaryl group where both of the aryl groups are heteroaryl groups. Exemplary heteroarylheteroaryl groups include, for example, 3-pyridylimidazole, 2-imidazolylpyrazine, and the like. Preferred optionally substituted heteroarylheteroaryl groups include: 2-(4-piperazinyl-3-pyridyl)furan, 20 diethyl(3-pyrazin-2-yl(4-pyridyl))amine, and dimethyl{2-[2-(5-methylpyrazin-2-yl)ethynyl](4-pyridyl)}amine.

"Optionally substituted" refers to the optional replacement of hydrogen with one or more monovalent or divalent radicals. Optionally substituted groups include those described herein, for each group in which a distinct definition for substitution is supplied. 25 Additionally, suitable substitution groups include, for example, hydroxyl, nitro, amino, imino, cyano, halo, thio, thioamido, amidino, imidino, oxo, oxamidino, methoxamidino, imidino, guanidino, sulfonamido, carboxyl, formyl, alkyl, substituted alkyl, haloloweralkyl, loweralkoxy, haloloweralkoxy, loweralkoxyalkyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, heteroarylcarbonyl, heteroaralkylcarbonyl, alkylthio, 30 aminoalkyl, cyanoalkyl, benzyl, pyridyl, pyrazolyl, pyrrole, thiophene, imidazolyl, and the like.

Representative substituted aminocarbonyl groups include, for example, those shown below. These can be further substituted by heterocyclyl groups and heteroaryl

groups as will be apparent to those having skill in the organic and medicinal chemistry arts in conjunction with the disclosure herein. Preferred aminocarbonyl groups include:

N-(2-cyanoethyl)carboxamide, N-(3-methoxypropyl)carboxamide,
 N-cyclopropylcarboxamide, N-(2-hydroxy-isopropyl)carboxamide, methyl
 5 2-carboxylamino-3-hydroxypropanoate, N-(2-hydroxypropyl)carboxamide,
 N-(2-hydroxy-isopropyl)carboxamide, N-[2-hydroxy-1-(hydroxymethyl)ethyl]carboxamide, N-(2-carboxylaminoethyl)acetamide, N-(2-(2-pyridyl)ethyl)carboxamide, N-(2-pyridylmethyl)carboxamide, N-(oxolan-2-ylmethyl)-
 10 carboxamide, N-(4-hydroxypyrrolidin-2-yl)carboxamide, N-[2-(2-hydroxyethoxy)ethyl]-carboxamide, N-(4-hydroxycyclohexyl)carboxamide, N-[2-(2-oxo-4-imidazolyl)ethyl]-carboxamide,
 N-(carboxylaminomethyl)acetamide, N-(3-pyrrolidinylpropyl)carboxamide, N-[1-(carboxylaminomethyl)pyrrolidin-3-yl]acetamide,
 N-(2-morpholin-4-ylethyl)carboxamide, N-[3-(2-oxopyrrolidinyl)propyl]carboxamide, 4-methyl-2-oxopiperazinecarbaldehyde, N-(2-hydroxy-3-pyrrolidinylpropyl)carboxamide,
 15 N-(2-hydroxy-3-morpholin-4-ylpropyl)carboxamide, N-{2-[(5-cyano-2-pyridyl)amino]ethyl}carboxamide, 3-(dimethylamino)pyrrolidinecarbaldehyde, N-[(5-methylpyrazin-2-yl)methyl]carboxamide, 2,2,2-trifluoro-N-(1-formylpyrrolidin-3-yl)acetamide,




Representative substituted alkoxy carbonyl groups include, for example, those shown below. These alkoxy carbonyl groups can be further substituted as will be apparent to those having skill in the organic and medicinal chemistry arts in conjunction with the disclosure herein.

25

Representative substituted alkoxy carbonyl groups include, for example, those shown below. These alkoxy carbonyl groups can be further substituted as will be apparent to those having skill in the organic and medicinal chemistry arts in conjunction with the disclosure herein.

5



The term "protected" with respect to hydroxyl groups, amine groups, and
 10 sulfhydryl groups refers to forms of these functionalities which are protected from
 undesirable reaction with a protecting group known to those skilled in the art such as
 those set forth in *Protective Groups in Organic Synthesis*, Greene, T.W.; Wuts, P. G. M.,
 John Wiley & Sons, New York, NY, (3rd Edition, 1999) which can be added or removed
 using the procedures set forth therein. Examples of protected hydroxyl groups include,
 15 but are not limited to, silyl ethers such as those obtained by reaction of a hydroxyl group
 with a reagent such as, but not limited to, t-butyldimethyl-chlorosilane,
 trimethylchlorosilane, triisopropylchlorosilane, triethylchlorosilane; substituted methyl
 and ethyl ethers such as, but not limited to methoxymethyl ether, methythiomethyl ether,
 benzyloxymethyl ether, t-butoxymethyl ether, 2-methoxyethoxymethyl ether,
 20 tetrahydropyranyl ethers, 1-ethoxyethyl ether, allyl ether, benzyl ether; esters such as, but
 not limited to, benzoylformate, formate, acetate, trichloroacetate, and trifluoroacetate.
 Examples of protected amine groups include, but are not limited to, amides such as,
 formamide, acetamide, trifluoroacetamide, and benzamide; imides, such as phthalimide,
 and dithiosuccinimide; and others. Examples of protected sulfhydryl groups include, but
 25 are not limited to, thioethers such as S-benzyl thioether, and S-4-picolyl thioether;

substituted S-methyl derivatives such as hemithio, dithio and aminothio acetals; and others.

A "pharmaceutically acceptable salt" includes a salt with an inorganic base, organic base, inorganic acid, organic acid, or basic or acidic amino acid. As salts of
5 inorganic bases, the invention includes, for example, alkali metals such as sodium or potassium; alkaline earth metals such as calcium and magnesium or aluminum; and ammonia. As salts of organic bases, the invention includes, for example, trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, and triethanolamine. As salts of inorganic acids, the instant invention includes, for example, hydrochloric acid,
10 hydroboric acid, nitric acid, sulfuric acid, and phosphoric acid. As salts of organic acids, the instant invention includes, for example, formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, and p-toluenesulfonic acid. As salts of basic amino acids, the instant invention includes, for example, arginine, lysine and ornithine.
15 Acidic amino acids include, for example, aspartic acid and glutamic acid.

As used herein, the term "pharmaceutically acceptable ester" refers to esters which hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly
20 alkanoic, alkenoic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Representative examples of particular esters include, but are not limited to, formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.

The term "pharmaceutically acceptable prodrugs" as used herein refers to those
25 prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term
30 "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., *Bioreversible*

Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

As used herein, "limit", "treat" and "treatment" are interchangeable terms as are "limiting" and "treating" and, as used herein, include preventative (e.g., prophylactic) and
5 palliative treatment or the act of providing preventative or palliative treatment.


"Treating" within the context of the instant invention, means an alleviation of symptoms associated with a disorder or disease, or halt of further progression or worsening of those symptoms, or prevention or prophylaxis of the disease or disorder. For example, within the context of treating patients in need of an inhibitor of PI3K, successful treatment
10 may include a reduction in the proliferation of capillaries feeding a tumor or diseased tissue, an alleviation of symptoms related to a cancerous growth or tumor, proliferation of capillaries, or diseased tissue, a halting in capillary proliferation, or a halting in the progression of a disease such as cancer or in the growth of cancerous cells. Treatment may also include administering the pharmaceutical formulations of the present invention in
15 combination with other therapies. For example, the compounds and pharmaceutical formulations of the present invention may be administered before, during, or after surgical procedure and/or radiation therapy. The compounds of the invention can also be administered in conjunction with other anti-cancer drugs including those used in antisense and gene therapy.

20 The PI3K inhibitors of this invention, as described herein, can be administered in the form of acid addition salts. The salts are conveniently formed by reacting a compound, if basic, with a suitable acid, such as have been described above. The salts are quickly formed in high yields at moderate temperatures, and often are prepared by merely isolating the compound from a suitable acidic wash as the final step of the
25 synthesis. The salt-forming acid is dissolved in an appropriate organic solvent, or aqueous organic solvent, such as an alkanol, ketone or ester. On the other hand, if the compound of this invention is desired in the free base form, it is isolated from a basic final wash step, according to the usual practice. A preferred technique for preparing hydrochlorides is to dissolve the free base in a suitable solvent and dry the solution thoroughly, as over
30 molecular sieves, before bubbling hydrogen chloride gas through it. It will also be recognized that it is possible to administer amorphous forms of the PI3K inhibitors.


The subject invention also includes isotopically-labeled PI3K inhibitors, which are structurally identical to those disclosed above, but for the fact that one or more atoms

are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. Compounds of the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds and of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically labeled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out known or referenced procedures and by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

Generally, the invention provides compounds having the formula I. The invention also provides tautomers of the compounds, pharmaceutically acceptable salts, esters and prodrugs of the compounds, and pharmaceutically acceptable salts, esters and prodrugs of the tautomers. Formula I has the following structure:



In one embodiment, W is:



wherein Z is selected from the group consisting of $-\text{CH}_2-$, $-\text{NH}-$, $-\text{O}-$, $-\text{S}-$, and $-\text{NR}_6-$, where R_6 is an alkyl or substituted alkyl group;

R_3 is absent or selected from the group consisting of alkyl, substituted alkyl, amino, alkylamino, aminoalkyl, dialkylamino, dialkylaminoalkyl, alkoxy, alkenyl, substituted alkenyl, alkynyl, carbonylamino, and alkoxycarbonyl; and

m and n are integers from 0-2;

X is a covalent bond or is selected from the group consisting of $-\text{CH}_2-$, $-\text{CHF}-$, $-\text{CF}_2-$, $-\text{NH}-$, $-\text{O}-$, $-\text{S}-$, and $-\text{NR}_5-$, where R_5 is an alkyl or substituted alkyl group;

Y is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;


R_1 is selected from the group consisting of hydrogen, halogen, carboxylic acid, and alkyl; and

R_2 is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

the tautomers thereof; and

the pharmaceutically acceptable salts, esters, or prodrugs thereof.

In another aspect of the compound of formula (I), X is a covalent bond and Y is a morpholinyl group as shown below:



wherein, R_3 , m, and n are as described above.

In another more particular embodiment of compound (I), X is a covalent bond and Y is an unsubstituted morpholinyl group.

In another more particular embodiment of compound (I), X is $-\text{NH}-$.

In another more particular embodiment of compound (I), Y is a heteroaryl or substituted heteroaryl group selected from pyridyl and alkoxy pyridyl.


In another more particular embodiment of compound (I), R_1 is hydrogen.

In another more particular embodiment of compound (I), R_2 is an aryl or substituted aryl group.

In another more particular embodiment of compound (I), R_2 is selected from the group consisting of phenyl, phenol, aniline, hydroxybenzyl, phenylalkoxycarbonyl, phenylcarbonylalkoxy, phenylaminocarbonyl, and phenylcarbonylamino.

In another more particular embodiment of compound (I), R_3 is absent.

5 In one aspect of the invention, a compound of formula (II) is provided:



wherein, X is selected from the group consisting of -NH-, -O-, and -S-;

Y is selected from the group consisting of heterocyclyl, substituted heterocyclyl,
10 aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

R_1 is hydrogen, halogen, or a carboxylic acid group;

R_2 is selected from the group consisting of aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

the tautomers thereof;

15 and the pharmaceutically acceptable salts, esters, or prodrugs thereof.

In another aspect of the compound of formula (II), X is -NH-.

In another aspect of the compound of formula (II), Y is a heteroaryl or substituted heteroaryl group selected from pyridyl and alkoxy pyridyl.


In another aspect of the compound of formula (II), R_1 is hydrogen.

20 In another aspect of the compound of formula (II), R_2 is an aryl or substituted aryl group.

In another aspect of the compound of formula (II), R_2 is selected from the group consisting of phenyl, phenol, aniline, hydroxybenzyl, phenylalkoxycarbonyl, phenylcarbonylalkoxy, phenylaminocarbonyl, and phenylcarbonylamino.

25 In another aspect of the compound of formula (II), R_3 is absent.

In another aspect of the invention, compounds of formula (XIII) are provided:



XIII

wherein,

X is selected from the group consisting of -NH-, -O-, and -S-;

5 Y is selected from the group consisting of heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, and substituted heteroaryl;

R₁ is hydrogen, halogen, or a carboxylic acid;

R₄ is independently selected from the group consisting of hydrogen, halogen, alkyl, substituted alkyl, hydroxy, alkoxy, amino, alkylamino, aminoalkyl, dialkylamino, dialkylaminoalkyl, aryl, heteroaryl, heterocyclyl, carbonylamino, and alkoxycarbonyl;

q is an integer from 1-5.

the tautomers thereof;


and the pharmaceutically acceptable salts, esters, or prodrugs thereof.

15 In another aspect of the compound of formula (XIII), X is -NH- and R₁ is hydrogen.

In another aspect of the compound of formula (XIII), R₄ is selected from the group consisting of hydrogen, halogen, alkyl, substituted alkyl, hydroxy, alkoxy, amino, alkylamino, aminoalkyl, dialkylamino, dialkylaminoalkyl, carbonylamino, and alkoxycarbonyl.

20 Other compounds of the invention are described above in the Summary of the Invention.

In other aspects, the present invention provides compositions that include the phosphatidylinositol 3-kinase inhibitor compounds described herein, and methods that utilize the phosphatidylinositol 3-kinase inhibitor compounds described herein. In addition to the compounds described above, the compositions and methods of the invention can also include and utilize compounds having the following formula:




its stereoisomers, tautomers, pharmaceutically acceptable salts, esters, and
 5 prodrugs, wherein

Y is selected from the groups consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- 10 (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- 15 (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- 20 (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- 25 (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- 30 (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- 5 (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

10 R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- 15 (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- 20 (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

25 Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z,
- (c) -S-,
- (d) -SO-,
- 30 (e) -SO₂-, and
- (f) -CH₂-,

wherein R^Z is H or substituted or unsubstituted alkyl group; and
 R^{4w} is selected from the group consisting of

- 5 (a) H,
(b) substituted or unsubstituted C_1 - C_6 -alkyl,
(c) $-COOR^{5w}$,
(d) $-CONH_2$,
(e) $-OR^{5w}$, and
(f) $-NHR^{5w}$,

wherein R^{5w} is H or C_1 - C_6 -alkyl; and

10 r is 0, 1, or 2.

In one aspect, the invention provides pharmaceutical formulation that include one or more compounds described herein in combination with a pharmaceutically acceptable carrier. The pharmaceutical formulations of the invention can include additional therapeutic agents including, for example, other conventional cytotoxic agents.
15 Representative other conventional cytotoxic agents include for example, irinotecan, topotecan, gemcitabine, gleevec, herceptin, 5-fluorouracil, leucovorin, carboplatin, cisplatin, taxanes, tezacitabine, cyclophosphamide, vinca alkaloids, imatinib, anthracyclines, rituximab, tamoxifen, CPT 11, and trastuzumab, and the like, and are considered to fall within the scope of this invention.

20 In another aspect, the invention provides methods for using the compounds described herein. For example, the compounds described herein can be used in the treatment of cancer. The compounds described herein can also be used in the manufacture of a medicament for the treatment of cancer.

The methods can utilize pharmaceutical formulations that include one or more
25 compounds described herein in combination with a pharmaceutically acceptable carrier. The methods can also utilize pharmaceutical formulations of the invention that include additional therapeutic agents.

In one embodiment, the invention provides a method for inhibiting phosphatidylinositol (PI) 3-kinase activity in a human or animal subject. In the method,
30 an amount of a compound described herein effective to inhibit phosphatidylinositol (PI)

3-kinase activity in the human or animal subject is administered to the human or animal subject.

In another embodiment, the invention provides a method of treating a patient in need of an inhibitor of phosphatidylinositol 3-kinase (PI3K) is provided. In the method, a pharmaceutical formulation containing an effective amount of a compound described herein is administered to a patient in need thereof.

In another embodiment, the invention provides a method for treating a condition (e.g., cancer) by modulation of phosphatidylinositol (PI) 3-kinase activity. In the method, an effective amount of a compound described herein is administered to a human or animal subject in need of such treatment.

In another embodiment, the invention provides a method for treating a cancer disorder in a human or animal subject. In the method, a composition comprising an amount of a compound described herein effective to treat cancer is administered to a human or animal subject in need thereof. As noted above, the administered composition can further include additional therapeutic agents (e.g., conventional cytotoxic agents).

In another embodiment, the invention provides a method for inhibiting tumor growth in a patient. In the method, an effective amount of a compound described herein is administered to a patient having a tumor.

In another embodiment, the invention provides a method for inhibiting the proliferation of capillaries in a patient. In the method, an effective amount of a compound described herein is administered to a patient in need.

The invention also provides methods of preparing pharmaceutical formulations comprising mixing any of the above-described compounds with a pharmaceutically acceptable carrier, water, or an aqueous solution.

PHARMACEUTICAL COMPOSITIONS

Pharmaceutical compositions of the present invention comprise a therapeutically effective amount of a phosphatidylinositol 3-kinase inhibitor compound described herein formulated together with one or more pharmaceutically acceptable carriers. As used herein, the term "pharmaceutically acceptable carrier" means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as lactose, glucose and sucrose; starches such as corn starch and

potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol; esters
5 such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming
10 agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator. The pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray, or a liquid aerosol or dry powder formulation
15 for inhalation.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers
20 such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants
25 such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a
30 sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally

employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

5 The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

10 In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form may be accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming
15 microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations may also be prepared by entrapping the drug in liposomes or
20 microemulsions which are compatible with body tissues.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature
25 and therefore melt in the rectum or vaginal cavity and release the active compound.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose,
30 glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate,

e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, acetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

10 The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner.

15 Examples of embedding compositions which can be used include polymeric substances and waxes.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

20 The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert

25 diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they

30 release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulations, ear drops, and the like are also contemplated as being within the scope of this invention.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Compositions of the invention may also be formulated for delivery as a liquid aerosol or inhalable dry powder. Liquid aerosol formulations may be nebulized predominantly into particle sizes that can be delivered to the terminal and respiratory bronchioles.

Aerosolized formulations of the invention may be delivered using an aerosol forming device, such as a jet, vibrating porous plate or ultrasonic nebulizer, preferably selected to allow the formation of an aerosol particles having with a mass medium average diameter predominantly between 1 to 5 μ . Further, the formulation preferably has balanced osmolarity ionic strength and chloride concentration, and the smallest aerosolizable volume able to deliver effective dose of the compounds of the invention to the site of the infection. Additionally, the aerosolized formulation preferably does not impair negatively the functionality of the airways and does not cause undesirable side effects.

Aerosolization devices suitable for administration of aerosol formulations of the invention include, for example, jet, vibrating porous plate, ultrasonic nebulizers and energized dry powder inhalers, that are able to nebulize the formulation of the invention into aerosol particle size predominantly in the size range from 1-5 μ . Predominantly in this application means that at least 70% but preferably more than 90% of all generated aerosol particles are within 1-5 μ range. A jet nebulizer works by air pressure to break a liquid solution into aerosol droplets. Vibrating porous plate nebulizers work by using a sonic vacuum produced by a rapidly vibrating porous plate to extrude a solvent droplet through a porous plate. An ultrasonic nebulizer works by a piezoelectric crystal that shears a liquid into small aerosol droplets. A variety of suitable devices are available,

including, for example, AeroNebTM and AeroDoseTM vibrating porous plate nebulizers (AeroGen, Inc., Sunnyvale, California), Sidestream[®] nebulizers (Medic-Aid Ltd., West Sussex, England), Pari LC[®] and Pari LC Star[®] jet nebulizers (Pari Respiratory Equipment, Inc., Richmond, Virginia), and AerosonicTM (DeVilbiss Medizinische Produkte (Deutschland) GmbH, Heiden, Germany) and UltraAire[®] (Omron Healthcare, Inc., Vernon Hills, Illinois) ultrasonic nebulizers.

Compounds of the invention may also be formulated for use as topical powders and sprays that can contain, in addition to the compounds of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants such as chlorofluorohydrocarbons.

Transdermal patches have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

According to the methods of treatment of the present invention, tumor growth is reduced or prevented in a patient such as a human or lower mammal by administering to the patient a therapeutically effective amount of a compound of the invention, in such amounts and for such time as is necessary to achieve the desired result. By a "therapeutically effective amount" of a compound of the invention is meant a sufficient amount of the compound to treat tumor growth, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

The total daily dose of the compounds of this invention administered to a human or other mammal in single or in divided doses can be in amounts, for example, from 0.01 to 50 mg/kg body weight or more usually from 0.1 to 25 mg/kg body weight. Single dose compositions may contain such amounts or submultiples thereof to make up the daily dose. In general, treatment regimens according to the present invention comprise administration to a patient in need of such treatment from about 10 mg to about 2000 mg of the compound(s) of this invention per day in single or multiple doses.

Methods of formulation are well known in the art and are disclosed, for example, in Remington: *The Science and Practice of Pharmacy*, Mack Publishing Company, Easton, Pa., 19th Edition (1995). Pharmaceutical compositions for use in the present invention can be in the form of sterile, non-pyrogenic liquid solutions or suspensions, coated capsules, suppositories, lyophilized powders, transdermal patches or other forms known in the art.

A "kit" as used in the instant application comprises a container for containing the pharmaceutical compositions and may also include divided containers such as a divided bottle or a divided foil packet. The container can be in any conventional shape or form as known in the art which is made of a pharmaceutically acceptable material, for example a paper or cardboard box, a glass or plastic bottle or jar, a resealable bag (for example, to hold a "refill" of tablets for placement into a different container), or a blister pack with individual doses for pressing out of the pack according to a therapeutic schedule. The container employed can depend on the exact dosage form involved, for example a conventional cardboard box would not generally be used to hold a liquid suspension. It is feasible that more than one container can be used together in a single package to market a single dosage form. For example, tablets may be contained in a bottle which is in turn contained within a box.

An example of such a kit is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process, recesses are formed in the plastic foil. The recesses have the size and shape of individual tablets or capsules to be packed or may have the size and shape to accommodate multiple tablets and/or capsules to be packed. Next, the tablets or capsules are placed in the recesses accordingly and the sheet of

relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are individually sealed or collectively sealed, as desired, in the recesses between the plastic foil and the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.

It maybe desirable to provide a written memory aid, where the written memory aid is of the type containing information and/or instructions for the physician, pharmacist or other health care provider, or subject, e.g., in the form of numbers next to the tablets or capsules whereby the numbers correspond with the days of the regimen which the tablets or capsules so specified should be ingested or a card which contains the same type of information. Another example of such a memory aid is a calendar printed on the card, e.g., as follows "First Week, Monday, Tuesday,"... etc ... "Second Week, Monday, Tuesday, ..." etc. Other variations of memory aids will be readily apparent. A "daily dose" can be a single tablet or capsule or several tablets or capsules to be taken on a given day. When the kit contains separate compositions, a daily dose of one or more compositions of the kit can consist of one tablet or capsule while a daily dose of another one or more compositions of the kit can consist of several tablets or capsules.

Another specific embodiment of a kit is a dispenser designed to dispense the daily doses one at a time in the order of their intended use. Preferably, the dispenser is equipped with a memory-aid, so as to further facilitate compliance with the regimen. An example of such a memory-aid is a mechanical counter, which indicates the number of daily doses that has been dispensed. Another example of such a memory-aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.

The kits of the present invention may also comprise, in addition to a PI3K inhibitor, one or more additional pharmaceutically active compounds. Preferably, the additional compound is another PI3K inhibitor or another compound useful to treat cancer, angiogenesis, or tumor growth. The additional compounds may be administered in the same dosage form as the PI3K inhibitor or in different dosage forms. Likewise, the

additional compounds can be administered at the same time as the PI3K inhibitor or at different times.

All references and patents cited herein are incorporated by reference.

The present invention, thus generally described, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.


EXAMPLES

Experimental

Compounds of the invention may be generally prepared using procedures well known to those skilled in the art, for example in accordance with the following representative methods (e.g., Methods 1 and 2 below) and reaction schemes.

METHOD 1

Resin bound 2-(pyrazolo)trihydropyrimidin-4-one (3) (Step 1)




15

Wang resin (1.0 g, 0.55 mmol, 1 eq) was suspended in toluene (10 mL) and DIEA (0.377 mL, 2.2 mmol, 4.0 eq) was added, followed by methyl malonyl chloride (0.236 mL, 2.2 mmol, 4.0 eq). The mixture was shaken overnight at room temperature. The resin was filtered and washed with CH₂Cl₂, MeOH, water, DMF, CH₂Cl₂ then dried to obtain resin bound methyl malonate 1. Resin 1 (300 mg, 0.165 mmol, 1.0 eq) was suspended in a solution of piperidine (16.3 μL, 0.165 mmol, 1.0 eq) and acetic acid (9.4 μL, 0.165 mmol, 1.0 eq) in DMF (3 mL) and the aldehyde (10.0 eq) was added. The mixture was shaken at room temperature overnight. The resin was filtered, washed with DMF and CH₂Cl₂, then dried to give the resin bound α,β unsaturated diester 2, which was used in the next step without analytics, since cleavage from the resin causes extensive decomposition. Resin 2 (300 mg, 0.165 mmol, 1.0 eq) was suspended in NMP (3 mL), and 1-H-pyrazole carboxamidine hydrochloride (121 mg 0.825 mmol, 5.0 eq) was added, followed by NaHCO₃ (35 mg, 0.412 mmol, 2.5 eq). The reaction mixture was shaken overnight at 50 °C, then the resin was filtered, washed with DMF, water, MeOH, CH₂Cl₂ and dried, to obtain the desired resin bound 6-R¹-4-oxo-2-pyrazolyl-

30

3,5,6-trihydropyrimidine-5-carboxylic acid 3. An analytical sample of the cleaved product was obtained treating the resin with 95% TFA/H₂O for 1.5 h at room temperature, filtering and evaporating under reduced pressure.


5 Resin bound 2-pyrazolopyrimidinone (4) (Step 2)



Step 2


Resin 3 (200 mg, 0.11 mmol, 1 eq) was suspended in 0.1 M solution of DDQ in toluene (2.5 mL, 253 mmol 2.3 eq) and the reaction mixture was shaken at 50°C overnight. The resin was filtered, washed with DMF, 20% aq AcOH, water, MeOH,
 10 CH₂Cl₂ and dried, to obtain the desired resin bound R¹-4-hydroxy-2-pyrazolylpyrimidine-5-carboxylic acid 4. An analytical sample of the cleaved product was obtained treating the resin with 95% TFA/H₂O for 1.5 h at room temperature, filtering and evaporating under reduced pressure.

15 PyBop Mediated Substitution with Amines in the 4 Position, (Step 3)




Step 3

A mixture of resin 4 (150 mg, 0.082 mmol, 1 eq), the amine of choice (10 eq), and PyBop (85 mg, 0.164 mmol, 2 eq) in NMP was shaken at room temperature overnight. The resin was filtered, washed with DMF, MeOH, and CH₂Cl₂, and dried, to obtain the
 20 desired resin bound 6-R¹-4-aminoalkyl (or aryl)-2-pyrazolylpyrimidine-5-carboxylic acid 5. An analytical sample of the cleaved product was obtained treating the resin 95% TFA/H₂O for 1.5 h at room temperature, filtering and evaporating under reduced pressure.

Sn_{Ar} with Morpholine in Position 2 (Step 4)**Step 4**

Resin 5 (100 mg, 0.055 mmol, 1 eq) was suspended in NMP, and morpholine (144 μ L, 144 mg, 1.65 mmol, 30 eq) was added, followed by acetic acid (31 μ L, 33 mg, 0.55 mmol, 10 eq). The reaction mixture was shaken at 90°C overnight. The resin was filtered and washed with DMF, water, MeOH, CH₂Cl₂, then dried. The resin was treated with 95% TFA/H₂O for 1.5 h at room temperature. Filtration and evaporation under reduced pressure afforded 6-R¹-4-alkyl (or aryl)amino-2-morpholino pyrimidine-5-carboxylic acid 6.

10

Decarboxylation (Step 5)**Step 5**

The carboxylic acid 6 was dissolved in a mixture of acetonitrile and water (1:1, 2 mL) and the solution was heated at 60 °C overnight. The solution was cooled down to room temperature and then lyophilized. After purification by reverse phase liquid chromatography, the desired trisubstituted pyrimidine 7 was obtained as a solid.

15

Example 1

(Synthesis of 3-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol)
6-(3-hydroxyphenyl)-4-oxo-2-pyrazolyl-3,5,6-trihydropyrimidine-5-carboxylic acid

20

Wang resin (1.0 g, 0.55 mmol, 1 eq) was suspended in toluene (10 mL) and DIEA (0.377 mL, 2.2 mmol, 4.0 eq) was added, followed by methyl malonyl chloride (0.236 mL, 2.2 mmol, 4.0 eq). The mixture was shaken overnight at room temperature. The resin was filtered and washed with CH₂Cl₂, MeOH, water, DMF, CH₂Cl₂ then dried to obtain resin bound methyl malonate (1). Resin 1 (300 mg, 0.165 mmol, 1.0 eq) was suspended in a solution of piperidine (16.3 μ L, 0.165 mmol, 1.0 eq) and acetic acid (9.4 μ L, 0.165 mmol, 1.0 eq) and 3-hydroxybenzaldehyde (201 mg, 1.65 mmol, 10.0 eq)

25

was added. The mixture was shaken at room temperature overnight. The resin was filtered, washed with DMF and CH₂Cl₂, dried, suspended in NMP, and 1-H-pyrazole carboxamide hydrochloride (121 mg 0.825 mmol, 5.0 eq) was added, followed by NaHCO₃ (35 mg, 0.412 mmol, 2.5 eq). The reaction mixture was shaken overnight at 50 °C, then the resin was filtered, washed with DMF, water, MeOH, CH₂Cl₂ and dried. To obtain an analytical sample, 20 mg of the resin were treated with 95% TFA/H₂O for 1.5 h at room temperature. Filtration and evaporation under reduced pressure afforded 6-(3-hydroxyphenyl)-4-oxo-2-pyrazolyl-3,5,6-trihydropyrimidine-5-carboxylic acid. HPLC (Buffer A: 0.1% TFA/H₂O; Buffer B: 0.1% TFA/CH₃CN; column: C18, 4.6x250mm; flow: 1mL/min; gradient: 2.1%, 5%-80% B in 36 min.): R_t= 14.70. LC/MS (ion spray, 50 eV, m/z): 275 (M+H₂O+H⁺).

4-hydroxy-6-(3-hydroxyphenyl)-2-pyrazolylpyrimidine-5-carboxylic acid

Resin bound 6-(3-hydroxyphenyl)-4-oxo-2-pyrazolyl-3,5,6-trihydropyrimidine-5-carboxylic acid (200 mg, 0.11 mmol, 1 eq) was suspended in 0.1 M solution of DDQ in toluene (2.5 mL, 253 mmol 2.3 eq) and the reaction mixture was shaken at 50 °C overnight. The resin was filtered, washed with DMF, 20% aq AcOH, water, MeOH, CH₂Cl₂ and dried. To obtain an analytical sample, 20 mg of the resin were treated with 95% TFA/H₂O for 1.5 h at room temperature. Filtration and evaporation under reduced pressure afforded 4-hydroxy-6-(3-hydroxyphenyl)-2-pyrazolylpyrimidine-5-carboxylic acid. HPLC (Buffer A: 0.1% TFA/H₂O; Buffer B: 0.1% TFA/CH₃CN; column: C18, 4.6x250mm; flow: 1mL/min; gradient: 2.1%, 5%-80% B in 36 min.): R_t= 15.78. LC-MS (ion spray, 50 eV, m/z): 299 (M+H⁺).

Resin bound 6-(3-hydroxyphenyl)-4-(1H-indazol-5-ylamino)

-2-pyrazolylpyrimidine-5-carboxylic acid

A mixture of resin bound 4-hydroxy-6-(3-hydroxyphenyl)-2-pyrazolylpyrimidine-5-carboxylic acid (150 mg, 0.082 mmol, 1 eq), 5-aminoindazole (110 mg, 0.82 mmol, 10 eq), and PyBop (85 mg, 0.164 mmol, 2 eq) in NMP was shaken at room temperature overnight. The resin was filtered, washed with DMF, MeOH, and CH₂Cl₂, and dried. To obtain an analytical sample, 20 mg of the resin were treated with 95% TFA/H₂O for 1.5 h at room temperature. Filtration and evaporation under reduced pressure afforded 6-(3-hydroxyphenyl)-4-(1H-indazol-5-ylamino)-2-pyrazolylpyrimidine-5-carboxylic acid.

HPLC (Buffer A: 0.1% TFA/H₂O; Buffer B: 0.1% TFA/CH₃CN; column: C18, 4.6x250mm; flow: 1mL/min; gradient: 2.1%, 5%-80% B in 36 min.): R_t= 20.72.

LC-MS (ion spray, 50 eV, m/z): 414 (M+H⁺).

Resin bound 6-(3-hydroxyphenyl)-4-(1H-indazol-5-ylamino)

5 -2-morpholin-4-ylpyrimidine-5-carboxylic acid

Resin bound 6-(3-hydroxyphenyl)-4-(1H-indazol-5-ylamino)-2-pyrazolylpyrimidine-5-carboxylic acid (100 mg, 0.055 mmol, 1 eq) was suspended in NMP, and morpholine (144 μL, 144 mg, 1.65 mmol, 30 eq) was added, followed by acetic acid (31 μL, 33mg, 0.55 mmol, 10 eq). The reaction mixture was shaken at 90 °C
10 overnight. The resin was filtered and washed with DMF, water, MeOH, CH₂Cl₂, then dried. The resin was treated with 95% TFA/H₂O for 1.5 h at room temperature. Filtration and evaporation under reduced pressure afforded 6-(3-hydroxyphenyl)-4-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid.

HPLC (Buffer A: 0.1% TFA/H₂O; Buffer B: 0.1% TFA/CH₃CN; column: C18, 4.6x250mm; flow: 1mL/min; gradient: 2.1%, 5%-80% B in 36 min.): R_t= 16.97.
15

LC-MS (ion spray, 50 eV, m/z): 433 (M+H⁺).

3-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol

6-(3-hydroxyphenyl)-4-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid was dissolved in a mixture of acetonitrile and water (1:1, 2 mL) and the
20 solution was heated at 60 °C overnight. The solution was cooled down to room temperature and then lyophilized. After purification by reverse phase liquid chromatography (Buffer A: 0.1% TFA/H₂O; Buffer B: 0.1% TFA/CH₃CN, column: C18, 5μ, 10x50mm, gradient 5%B-95%B in 9 min) the Bis TFA salt of 3-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol was obtained as a pale yellow solid.


25 ¹H-NMR (HCl salt, 60% CD₃CN/D₂O, 300 MHz): 8.09 (s, 1H), 8.03 (bs, 1H), 7.61 (1H, d, J= 8.7), 7.55 (bm, 1H), 7.38 (app. t, 1H, J= 7.8), 7.17 (bd, 1H, J= 7.8), 7.10 (bs, 1H), 7.06 (d, 1H, J= 8.7), 6.42 (bs, 1H), 3.75 (app. s, 8H).

HPLC (Buffer A: 0.1% TFA/H₂O; Buffer B: 0.1% TFA/CH₃CN; column: C18, 4.6x250mm; flow: 1mL/min; gradient: 2.1%, 5%-80% B in 36 min.): R_t= 18.17.


30 LC-MS (ion spray, 50 eV, m/z): 389 (M+H⁺).

METHOD 2

Solution Phase Synthesis of 3-[2-Morpholin-4-yl-6-(3-pyridylamino)pyrimidin-4-yl]phenol



Step 1:




To a stirred solution of 3'-hydroxyacetophenone (1eq) and benzyl bromide (1.5 eq) in dry DMF under N₂, solid K₂CO₃ (2 eq) was added in one portion. The reaction mixture was stirred at 60° C for 3 days, then cooled down to room temperature. Most of the DMF was distilled off under reduced pressure. The residue was taken up in EtOAc and washed with 1N HCl, H₂O, Brine and dried (Na₂SO₄). Evaporation of the solvent under reduced pressure afforded a brown oil which was about a 1:1 mixture of the starting material and the desired product. The latter was isolated by chromatography on silica gel (EtOAc/Hexanes, 1:1) affording the desired 3'-benzyloxy acetophenone (51%). See for example: Schmidhammer, H.; Brossi, A. *J. Org. Chem.* 1983, 48, 1469.

TLC (silica gel, Ethyl acetate/hexanes 1:2, vanillin stain): R_f = 0.58, orange brown (R_f starting material = 0.28).

¹H NMR (CDCl₃, 300MHz): 7.6-7.1 (9H, m), 5.11 (2H, s, CH₂Ph); 2.59 (3H, s, CH₃).

Step 2:




A round bottom flask, oven dried and kept under N₂ atmosphere, was charged
 5 with potassium *tert*-butoxide (2.2 eq) and dry toluene was added. The suspension was
 cooled down to 0 °C and a solution of 3'-benzyloxy acetophenone (1eq) and
 diethylcarbonate (2 eq) in toluene was added dropwise *via* a dropping funnel, with
 vigorous stirring. During the addition the temperature should not raise above 10 °C.
 After the end of the addition the reaction mixture was stirred at room temperature for 1h
 10 and then at 60 °C overnight. The reaction mixture was again cooled down to room
 temperature and quenched with a 1:10 mixture of acetic acid and water. The addition
 must be slow and occasional cooling might be necessary to keep the temperature below
 20 °C. The two phases were separated and the aqueous phase was extracted with EtOAc
 (x3). The organic extracts were collected and dried (Na₂SO₄). After evaporation of the
 15 solvent under reduced pressure of crude ethyl 3-oxo-3-[3-(phenylmethoxy)phenyl]propanoate were obtained. The compound could be carried on to
 the next step without further purification.

TLC (silica gel, ethyl acetate/hexanes 1:5, vanillin stain): R_f = 0.26, faint orange
 brown

20 LC-MS (ion spray, 50 eV, m/z): 299 (M+ H⁺).

¹H NMR (CDCl₃, 300MHz): 7.6-7.1 (9H, m); 5.10 (2H, bs, CH₂Ph); 4.21 (2H, q,
 J = 7.2 Hz OCH₂); 3.96 (2H, s, COCH₂); 1.25 (3H, t, J = 7.2 Hz, CH₃).

Step 3:



25

In a round bottom flask, oven dried and kept under N₂ atmosphere, Cs₂CO₃
 (1.5 eq) was suspended in dry DMF. Morpholino formamidine hydrobromide (1.2 eq)
 was added, followed by ethyl 3-oxo-3-[3-(phenylmethoxy)phenyl] propanoate (1eq). The


reaction mixture was stirred at 115 °C overnight, then cooled down to room temperature. The DMF was distilled off under reduced pressure and the residue was taken up in water, neutralizing with 5% HCl solution. The aqueous phase was then extracted with CH₂Cl₂ (x5). The organic extracts were collected and dried (Na₂SO₄). After evaporation of the solvent under reduced pressure the desired 2-morpholin-4-yl-6-[3-(phenylmethoxy)phenyl]-3-hydropyrimidin-4-one were obtained as an off white solid (60%). The crude is already pure enough for the next step, but it can be purified further by trituration with acetonitrile.

TLC (silica gel, CH₂Cl₂/MeOH 1:10): R_f = 0.32 (R_f of the starting material = 0.9).

LC-MS (ion spray, 50 eV, m/z): 364 (M+ H⁺).

¹H NMR (CDCl₃, 300MHz): 7.65-7.3 (8H, m); 7.06 (1H, ddd, J = 8.4, 2.7, 0.9 Hz); 6.25 (1H, s); 5.13 (2H, s, CH₂Ph); 3.83 (8H, bs, morpholine).

Step 4:




2-morpholin-4-yl-6-[3-(phenylmethoxy)phenyl]-3-hydropyrimidin-4-one (1eq) was dissolved in CH₂Cl₂ in a round bottom flask, oven dried and kept under N₂ atmosphere. The compound is not completely soluble. Triethylamine was added (1.4 eq) followed by N-Phenyl trifluoromethanesulfonamide (1.2 eq) and DMAP (10 mol%). The reaction mixture was stirred at room temperature overnight, obtaining a bright orange solution. The solvent was evaporated under reduced pressure and the residue purified by chromatography on silica gel (ethyl acetate/hexanes 1:5), obtaining (99%) the desired 2-morpholin-4-yl-6-[3-(phenylmethoxy)phenyl]pyrimidin-4-yl (trifluoromethyl) sulfonate.

TLC (silica gel, EtOAc/Hexanes 1:5): R_f = 0.31.


^1H NMR (CDCl_3 , 300MHz): 7.64 (1H, dd, J =2.4, 1.5 Hz); 7.55 (1H, app. dt, J =7.8, 1.2 Hz); 7.2-7.3 (6H, m); 7.12 (1H, ddd, J = 8.4, 2.4, 0.9 Hz); 6.66 (1H, s, pyrimidine CH), 5.14 (2H, s, CH_2Ph), 3.86 (4H, bm, morpholine); 3.79 (4H, m, morpholine).

Step 5:



A round bottom flask, oven dried and kept under N_2 atmosphere was charged with Cs_2CO_3 (1.4 eq), $\text{Pd}(\text{OAc})_2$ (5 mol%), and S-(-)-BINAP (1.5 x mol of Pd catalyst). The flask was purged with N_2 for about 5-10 min and a solution of 2-morpholin-4-yl-6-[3-(phenylmethoxy)phenyl]pyrimidin-4-yl (trifluoromethyl) sulfonate (1eq) in dry THF (20 mL) was added via a syringe, followed by 3-aminopyridine (2 eq.) in one portion. The flask was equipped with a reflux condenser, purged again with N_2 for 5 min and the reaction mixture was refluxed overnight. An efficient stirring is very important. The reaction mixture was cooled down to room temperature and the solvent was evaporated under reduced pressure. The residue was washed with water (x2) and triturated with methanol to afford the desired {2-morpholin-4-yl-6-[3-(phenylmethoxy)phenyl]pyrimidin-4-yl}-3-pyridylamine.

Step 6:



{2-Morpholin-4-yl-6-[3-(phenylmethoxy)phenyl]pyrimidin-4-yl}-3-pyridylamine (1 eq) is suspended in ethanol in a round bottom flask, purged with N_2 . 10% Pd/C (20% wt) was added. The flask was evacuated and filled up with H_2 (contained in a balloon) for five times, then the reaction mixture was stirred under H_2 for 20 h. The catalyst was filtered off through a pad of celite washing thoroughly with EtOH, MeOH, CH_2Cl_2 , and

acetonitrile (almost one liter of the mixture of solvents was used to ensure the complete solubilization of the product). The solvent was evaporated under reduced pressure and the residue purified by reverse phase chromatography (Buffer A: 0.1% TFA in H₂O, Buffer B: 0.1% TFA in CH₃CN; Column: Waters, C18, 47x300 mm; gradient: 1.1%,
5 10%-60%B in 45 min). The free base thus obtained was lyophilized from a 1:1 mixture of acetonitrile and 1N HCl, obtaining the desired 3-[2-morpholin-4-yl-6-(3-pyridylamino)pyrimidin-4-yl]phenol as the bis HCl salt. The spectral data are the following:







HPLC: (Buffer A: 0.1% TFA in H₂O, Buffer B: 0.1% TFA in CH₃CN; Column:
10 Waters, C18, 4.6x250 mm; gradient: 4.2%, 5%-80%B in 18 min) R_t= 4.47.







LC-MS (ion spray, 50 eV, m/z): 350 (M+ H⁺).

¹H NMR (DMSO+D₂O, 300MHz): 9.22 (1H, bs), 8.37 (2H, app d, J= 5.7), 7.79 (1H, dd, J= 7.2, 5.4), 7.43, (2H, m). 7.30 (1H, app t, J= 7.5), 6.89 (1H, dd, J= 7.0, 2.1), 6.59 (1H, s), 3.6-3.8 (8H, m).








15 Compounds of the following Examples were synthesized following the synthetic method described above in Methods 1 and 2. The precursors are readily recognizable by one skilled in the art and are commercially available from Aldrich (Milwaukee, WI), Acros Organics (Pittsburgh, PA), Biosynth International (Naperville, IL), Asymchem International, Inc. (Durham, NC) Maybridge Chemical Company Ltd. (Cornwall), and/or
20 UK Peakdale Molecular (High Peak, UK).








The compounds were named using ACD/Name v. 5.04, 2001 and Nomenclator (v. 6.0) from ChemInnovation Software, Inc.








Example	Structure	Name	LC/MS m/z (MH ⁺)
2		N-[6-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-6-amine	431.5
3		4-(3-hydroxyphenyl)-6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	433.4
4		4-[3-(2-hydroxyethoxy)phenyl]-6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	477.5
5		4-(1H-indazol-5-ylamino)-2-morpholin-4-yl-6-(4-phenoxyphenyl)pyrimidine-5-carboxylic acid	509.5
6		4-(2,3-dihydro-1,4-benzodioxin-6-yl)-6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	475.5
7		4-(1H-indazol-5-ylamino)-6-[4-(methylsulfonyl)phenyl]-2-morpholin-4-ylpyrimidine-5-carboxylic acid	495.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
8		4-[3-(4-tert-butylphenoxy)phenyl]-6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	565.6
9		4-[3-(3,5-dichlorophenoxy)phenyl]-6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	578.4
10		4-(4-tert-butylphenyl)-6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	473.5
11		4-(1H-indazol-5-ylamino)-2-morpholin-4-yl-6-phenylpyrimidine-5-carboxylic acid	417.4
11a			
12		N-[6-(4-methoxy-3-methylphenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	417.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
13		2-{3-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenoxy}ethanol	433.5
14		N-[2-morpholin-4-yl-6-(4-phenoxyphenyl)pyrimidin-4-yl]-1H-indazol-5-amine	465.5
15		N-{6-[4-(methylsulfonyl)phenyl]-2-morpholin-4-ylpyrimidin-4-yl}-1H-indazol-5-amine	451.5
16		N-{6-[3-(4-tert-butylphenoxy)phenyl]-2-morpholin-4-ylpyrimidin-4-yl}-1H-indazol-5-amine	521.6
17		N-{6-[3-(3,5-dichlorophenoxy)phenyl]-2-morpholin-4-ylpyrimidin-4-yl}-1H-indazol-5-amine	534.4
18		N-[6-(4-tert-butylphenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	429.5
19		N-(2-morpholin-4-yl-6-phenylpyrimidin-4-yl)-1H-indazol-5-amine	373.4

Example	Structure	Name	LC/MS m/z (MH ⁺)
20		4-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	389.4
21		N-[6-(3-fluorophenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	391.4
22		N-[6-(4-fluorophenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	391.4
23		N-[6-(2-fluorophenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	391.4
24		N-[6-(3-chlorophenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	407.9
25		N-[2-morpholin-4-yl-6-(3-nitrophenyl)pyrimidin-4-yl]-1H-indazol-5-amine	418.4
26		N-[2-morpholin-4-yl-6-[3-(trifluoromethoxy)phenyl]pyrimidin-4-yl]-1H-indazol-5-amine	457.4








Example	Structure	Name	LC/MS m/z (MH ⁺)
27		N-{2-morpholin-4-yl-6-[3-(trifluoromethyl)phenyl]pyrimidin-4-yl}-1H-indazol-5-amine	441.4
28		N-{6-[3-(benzyloxy)phenyl]-2-morpholin-4-ylpyrimidin-4-yl}-1H-indazol-5-amine	479.6
29		N-{6-(3-ethoxyphenyl)-2-morpholin-4-ylpyrimidin-4-yl}-1H-indazol-5-amine	417.5
30		3-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]benzonitrile	398.4
31		N-{6-(3-methylphenyl)-2-morpholin-4-ylpyrimidin-4-yl}-1H-indazol-5-amine	387.5
32		ethyl 4-[4-(3-hydroxyphenyl)-6-(1H-indazol-5-ylamino)pyrimidin-2-yl]piperazine-1-carboxylate	460.5
33		3-[2-(4-acetylpiperazin-1-yl)-6-(1H-indazol-5-ylamino)pyrimidin-4-yl]phenol	430.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
34		3-{6-[(1-acetyl-2,3-dihydro-1H-indol-6-yl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	432.5
35		3-[6-(2,3-dihydro-1H-inden-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	389.5
36		3-[6-(9H-fluoren-2-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	437.5
37		3-[6-(2,3-dihydro-1,4-benzodioxin-6-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	407.4
38		3-{6-[(3,4-dimethoxyphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	409.5
39		3-[6-(2,3-dihydro-1H-indol-6-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	390.5
40		3-[6-(1H-indazol-6-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	389.4

Example	Structure	Name	LC/MS m/z (MH ⁺)
41		3-[6-(1,3-benzodioxol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	393.4
42		3-[6-[(3-chloro-4-methoxyphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl]phenol	413.9
43		5-[[6-(3-hydroxyphenyl)-2-morpholin-4-ylpyrimidin-4-yl]amino]-2-methoxyphenol	395.4
44		3-[6-[(3-fluoro-4-methoxyphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl]phenol	397.4
45		5-[[6-(3-hydroxyphenyl)-2-morpholin-4-ylpyrimidin-4-yl]amino]-1,3-dihydro-2H-benzimidazol-2-one	405.4
46		3-[6-[(3,4-dimethylphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl]phenol	377.5
47		3-(2,6-dimorpholin-4-ylpyrimidin-4-yl)phenol	343.4








Example	Structure	Name	LC/MS m/z (MH ⁺)
48		4-[[6-(3-hydroxyphenyl)-2-morpholin-4-yl]amino]-2-nitrophenol	410.4
49		2-chloro-4-[[6-(3-hydroxyphenyl)-2-morpholin-4-yl]amino]phenol	399.8
50		3-{6-(1H-indazol-5-ylamino)-2-[(2-methoxyethyl)amino]pyrimidin-4-yl}phenol	377.4
51		3-[2-azepan-1-yl-6-(1H-indazol-5-ylamino)pyrimidin-4-yl]phenol	401.5
52		3-[2-(1,4-diazepan-1-yl)-6-(1H-indazol-5-ylamino)pyrimidin-4-yl]phenol	402.5
53		3-[2-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-6-(1H-indazol-5-ylamino)pyrimidin-4-yl]phenol	417.5
54		3-[6-(1H-indazol-5-ylamino)-2-thiomorpholin-4-yl]phenol	405.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
55		N-[6-(3-methoxyphenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	403.5
56		3-{6-[(4-methylbenzyl)(pyridin-2-ylmethyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	468.6
57		3-{2-morpholin-4-yl-6-[(2-pyridin-4-ylethyl)amino]pyrimidin-4-yl}phenol	378.4
58		3-{6-[(6-methoxypyridin-3-yl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	380.4
59		3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	350.4
60		3-[6-(dibenzylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	453.6
61		3-{6-[benzyl(1,3-thiazol-2-ylmethyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	460.6

Example	Structure	Name	LC/MS m/z (MH ⁺)
62		3-(2-morpholin-4-yl-6- {[(1R)-1- phenylethyl]amino}pyrimidin -4-yl)phenol	377.5
63		3-(6-anilino-2-morpholin-4- ylpyrimidin-4-yl)phenol	349.4
64		3-{2-morpholin-4-yl-6- [(3,4,5- trimethoxyphenyl)amino]pyri midin-4-yl}phenol	439.5
65		3-{6-[(4- butoxyphenyl)amino]-2- morpholin-4-ylpyrimidin-4- yl}phenol	421.5
66		3-(2-morpholin-4-yl-6-{[4- (pentyloxy)phenyl]amino}py rimidin-4-yl)phenol	435.5
67		3-(6-{[4- (hexyloxy)phenyl]amino}-2- morpholin-4-ylpyrimidin-4- yl)phenol	449.6
68		3-[6-(1H-benzimidazol-6- ylamino)-2-morpholin-4- ylpyrimidin-4-yl]phenol	389.4

Example	Structure	Name	LC/MS m/z (MH ⁺)
69		4-[4-(3-hydroxyphenyl)-6-(1H-indazol-5-ylamino)pyrimidin-2-yl]piperazine-1-carbaldehyde	416.5
70		methyl 3-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]benzoate	431.5
71		4-[4-(3-methoxyphenyl)-6-morpholin-4-ylpyrimidin-2-yl]morpholine	357.4
72		2-[6-(1H-indazol-5-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	389.4
73		3-[6-[(2-methoxyethyl)amino]-2-morpholin-4-ylpyrimidin-4-yl]phenol	331.4
74		2-ethyl-2-[[6-(3-hydroxyphenyl)-2-morpholin-4-ylpyrimidin-4-yl]amino]propane-1,3-diol	375.4
75		3-[6-(methylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	287.3








Example	Structure	Name	LC/MS m/z (MH ⁺)
76		3-{6-[2-(hydroxymethyl)pyrrolidin-1-yl]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	357.4
77		3-{6-[(3-aminocyclohexyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	370.5
78		3-{6-[(1R,2R)-2-aminocyclohexyl]amino}-2-morpholin-4-yl}pyrimidin-4-yl}phenol	370.5
79		3-{6-[(4-hydroxycyclohexyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	371.4
80		1-[6-(3-hydroxyphenyl)-2-morpholin-4-yl}pyrimidin-4-yl}piperidin-4-ol	357.4
81		3-{6-[(3R,5S)-3,5-dimethylmorpholin-4-yl]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	371.4
82		3-{2-morpholin-4-yl-6-[4-(4-nitrophenyl)piperazin-1-yl]}pyrimidin-4-yl}phenol	463.5








Example	Structure	Name	LC/MS m/z (MH ⁺)
83		3-[6-[4-(3-chlorophenyl)piperazin-1-yl]-2-morpholin-4-yl]pyrimidin-4-yl}phenol	453.0
84		3-[6-[4-(1,3-benzodioxol-5-ylmethyl)piperazin-1-yl]-2-morpholin-4-yl]pyrimidin-4-yl}phenol	476.5
85		3-[2-morpholin-4-yl-6-(4-pyridin-2-yl)piperazin-1-yl]pyrimidin-4-yl}phenol	419.5
86		3-[6-(4-acetylpiperazin-1-yl)-2-morpholin-4-yl]pyrimidin-4-yl}phenol	384.4
87		3-[6-(1,4-diazepan-1-yl)-2-morpholin-4-yl]pyrimidin-4-yl}phenol	356.4
88		3-[6-(4-methyl-1,4-diazepan-1-yl)-2-morpholin-4-yl]pyrimidin-4-yl}phenol	370.5
89		3-[2-morpholin-4-yl-6-[(pyridin-2-ylmethyl)amino]pyrimidin-4-yl}phenol	364.4

Example	Structure	Name	LC/MS m/z (MH ⁺)
90		3-{2-morpholin-4-yl-6-[(pyridin-3-ylmethyl)amino]pyrimidin-4-yl}phenol	364.4
91		3-{2-morpholin-4-yl-6-[(pyridin-4-ylmethyl)amino]pyrimidin-4-yl}phenol	364.4
92		3-{2-morpholin-4-yl-6-[(2-pyridin-2-ylethyl)amino]pyrimidin-4-yl}phenol	378.4
93		3-{2-morpholin-4-yl-6-[(2-pyridin-3-ylethyl)amino]pyrimidin-4-yl}phenol	378.4
94		3-(6-{[3-(1H-imidazol-1-yl)propyl]amino}-2-morpholin-4-ylpyrimidin-4-yl)phenol	381.4
95		3-{6-[(4-methylbenzyl)(pyridin-3-ylmethyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	468.6
96		3-(6-{[bis(2,4-dimethylphenyl)methyl]amino}-2-morpholin-4-ylpyrimidin-4-yl)phenol	495.6

Example	Structure	Name	LC/MS m/z (MH ⁺)
97		3-{6-[(2-methoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	379.4
98		3-{6-[(3-methoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	379.4
99		3-{6-[(4-methoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	379.4
100		3-{6-[(2,4-dimethoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	409.5
101		3-{6-[(2,5-dimethoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	409.5
102		3-{6-[(2,3-dimethoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	409.5
103		3-{6-[(2-ethoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	393.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
104		3-{6-[(4-ethoxyphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	393.5
105		3-{6-[(2,5-diethoxyphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	437.5
106		3-{6-[(2-methoxy-6-methylphenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	393.5
107		3-{2-morpholin-4-yl-6-[(3-phenoxyphenyl)amino]pyrimidin-4-yl}phenol	441.5
108		3-{2-morpholin-4-yl-6-[(4-phenoxyphenyl)amino]pyrimidin-4-yl}phenol	441.5
109		3-{6-[(3-(benzyloxy)phenyl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	455.5
110		3-{6-[(4-methoxydibenzo[b,d]furan-3-yl)amino]-2-morpholin-4-ylpyrimidin-4-yl}phenol	469.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
111		2-[[6-(3-hydroxyphenyl)-2-morpholin-4-yl]pyrimidin-4-yl]amino}phenol	365.4
112		3-{6-[(3-hydroxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	365.4
113		3-{6-[(4-hydroxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	365.4
114		4-chloro-2-[[6-(3-hydroxyphenyl)-2-morpholin-4-yl]pyrimidin-4-yl]amino}phenol	399.8
115		3-{[6-(3-hydroxyphenyl)-2-morpholin-4-yl]pyrimidin-4-yl}amino}-1,1'-biphenyl-4-ol	441.5
116		3-{6-[(4-anilino-2-methoxyphenyl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	470.5
117		3-{6-[(1-ethyl-2-methyl-1H-benzimidazol-5-yl)amino]-2-morpholin-4-yl}pyrimidin-4-yl}phenol	431.5







Example	Structure	Name	LC/MS m/z (MH ⁺)
118		N-(4-ethoxy-3-[[6-(3-hydroxyphenyl)-2-morpholin-4-yl]pyrimidin-4-yl]amino)phenyl)acetamide	450.5
119		3-[6-(1H-1,2,3-benzotriazol-6-ylamino)-2-morpholin-4-yl]pyrimidin-4-yl]phenol	390.4
120		2-methoxy-5-[(2-morpholin-4-yl-6-phenylpyrimidin-4-yl)amino]phenol	379.4
121		3-(6-amino-2-morpholin-4-yl)pyrimidin-4-yl]phenol	273.3
122		N-{2-morpholin-4-yl-6-[3-(2-piperidin-1-ylethoxy)phenyl]pyrimidin-4-yl}-1H-indazol-5-amine	500.6
123		4-(3-methoxyphenyl)-2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidine-5-carboxylic acid	408.4
124		3-{6-[4-(3-methoxyphenyl)piperazin-1-yl]-2-morpholin-4-yl}pyrimidin-4-yl]phenol	448.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
125		3-{2-morpholin-4-yl-6-[4-(2-morpholin-4-yl-2-oxoethyl)piperazin-1-yl]pyrimidin-4-yl}phenol	469.6
126		3-{2-morpholin-4-yl-6-[4-(1-phenylethyl)piperazin-1-yl]pyrimidin-4-yl}phenol	446.6
127		3-{2-morpholin-4-yl-6-[4-(2-phenylethyl)piperazin-1-yl]pyrimidin-4-yl}phenol	446.6
128		3-(6-{4-[2-(dimethylamino)ethyl]piperazin-1-yl}-2-morpholin-4-ylpyrimidin-4-yl)phenol	413.5
129		3-[6-(3,4-dihydro-2H-1,5-benzodioxepin-7-ylamino)-2-morpholin-4-ylpyrimidin-4-yl]phenol	421.5
130		3-(6-([3-(cyclopentyloxy)-4-methoxyphenyl]amino)-2-morpholin-4-ylpyrimidin-4-yl)phenol	463.5
131		3-[6-(1H-indazol-5-ylamino)-2-(1-oxidothiomorpholin-4-yl)pyrimidin-4-yl]phenol	421.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
132		3-[2-(2,6-dimethylmorpholin-4-yl)-6-(1H-indazol-5-ylamino)pyrimidin-4-yl]phenol	417.5
133		5-[[6-(3-hydroxyphenyl)-2-morpholin-4-yl]pyrimidin-4-yl]amino}pyridin-2-ol	366.4
134		6-(3-fluorophenyl)-2-morpholin-4-yl-N-pyridin-3-ylpyrimidin-4-amine	352.4
135		2-morpholin-4-yl-N-pyridin-3-yl-6-[3-(trifluoromethyl)phenyl]pyrimidin-4-amine	402.4
136		6-(3-methoxyphenyl)-2-morpholin-4-yl-N-pyridin-3-ylpyrimidin-4-amine	364.4
137		3-[2-morpholin-4-yl-6-(pyrimidin-2-ylamino)pyrimidin-4-yl]phenol	351.4
138		3-[2-morpholin-4-yl-6-(pyrazin-2-ylamino)pyrimidin-4-yl]phenol	351.4

Example	Structure	Name	LC/MS m/z (MH ⁺)
139		3-[6-(isoquinolin-5-ylamino)-2-morpholin-4-yl]pyrimidin-4-yl]phenol	400.5
140		3-[2-morpholin-4-yl-6-(quinolin-6-ylamino)pyrimidin-4-yl]phenol	400.5
141		3-[2-morpholin-4-yl-6-(quinolin-3-ylamino)pyrimidin-4-yl]phenol	400.5
142		3-[2-morpholin-4-yl-6-(pyridin-2-ylamino)pyrimidin-4-yl]phenol	350.4
143		3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl butyrate	420.5
144		3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl acetate	392.4
145		3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl pivalate	434.5

Example	Structure	Name	LC/MS m/z (MH ⁺)
146		3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl 2-methylpropanoate	420.5
147		6-(3-aminophenyl)-2-morpholin-4-yl-N-pyridin-3-ylpyrimidin-4-amine	349.4
148		2-fluoro-3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	368.4
149		3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl valinate	449.5
150		2-chloro-5-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	384.8
151		N-[3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl]methanesulfonamide	427.5
152		4-fluoro-3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	368.4

Example	Structure	Name	LC/MS m/z (MH ⁺)
153		4-bromo-3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	429.3
154		2-methyl-5-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	364.4
155		methyl 3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenyl carbonate	408.4
156		4-methyl-3-[2-morpholin-4-yl-6-(pyridin-3-ylamino)pyrimidin-4-yl]phenol	364.4
157		6-(3-hydroxyphenyl)-2-morpholin-4-ylpyrimidin-4-ol	274.3
158		4-(1H-indazol-5-ylamino)-6-(4-methoxy-3-methylphenyl)-2-morpholin-4-ylpyrimidine-5-carboxylic acid	461


Example	Structure	Name	LC/MS m/z (MH ⁺)
159		4-(1H-indazol-5-ylamino)-2-morpholin-4-yl-6-quinolin-3-ylpyrimidine-5-carboxylic acid	468
160		N-(2-morpholin-4-yl-6-quinolin-3-ylpyrimidin-4-yl)-1H-indazol-5-amine	424
161		N-[6-(3-bromophenyl)-2-morpholin-4-ylpyrimidin-4-yl]-1H-indazol-5-amine	452

Example 162

4-Substituted Pyrimidinyl Compounds

Modifications in the 4- position of the pyrimidinyl core can be accomplished starting from the 4- ester moiety, as shown in Scheme 1 below. The ester 1 can be reduced to the alcohol 2 and then reoxydized to the corresponding aldehyde 3. The aldehyde can be used as a substrate for reductive amination with primary or secondary amines to afford 4-alkyl (or dialkyl)aminomethyl substituted pyrimidines 4 (representative procedure given below). As an alternate route, alcohol 2 can be converted to a good leaving group such as mesylate, tosylate (5) triflate and the like, and reacted with suitable nucleophiles such as primary or secondary amines, alcohols, thiols. As an additional alternate route the ester can be hydrolyzed to the carboxylic acid, which can be

easily coupled with a variety of primary and secondary amines to afford 4-amides 6. Reduction of the amide will afford the desired 4- alkyl (or dialkyl)aminomethyl substituted pyrimidines 4 (see scheme 1). Compound 1 and its analogs can be obtained via Knoevenagel condensation of suitable aldehydes with ethyl acetoacetate, followed by oxidation of the dihydropyrimidine core with DDQ, with a procedure essentially identical to the one used in the solid phase synthesis of this class of compounds (other known agents e.g. CAN, can also be used in the aromatization step).



Scheme 1

General procedure for the synthesis of {[6-(3-methoxyphenyl)-2-morpholin-4-yl-4-(3-pyridylamino)pyrimidin-5-yl]methyl}dialkylamines.

[6-(3-methoxyphenyl)-2-morpholin-4-yl-4-(3-pyridylamino)pyrimidin-5-yl]formaldehyde (3)

Ester 1 is suspended in THF and DIBALH (1.6 N solution in THF, 3 eq) is added dropwise via a syringe. The reaction mixture is stirred at 50 °C overnight, then cooled down to room temperature and quenched with water. Product 2 precipitates and is filtered off, dried and used as is in the following step. Alcohol 2 is dissolved in DMA, and MnO₂ (xs) is added. The reaction mixture is stirred at room temperature overnight, and the solid is filtered off. The resulting clear solution is concentrated distilling off the solvent under reduced pressure and water is added. The precipitate thus obtained is

filtered off, triturated with more water and dried, obtaining aldehyde 3, which is not purified further.


{[6-(3-methoxyphenyl)-2-morpholin-4-yl-4-(3-pyridylamino)
pyrimidin-5-yl]methyl}dialkylamines (4)

- 5 A mixture of aldehyde 3 (1 eq), the desired amine (2.5-3 eq), and NaCNBH₃ in MeOH is refluxed overnight. The reaction mixture is cooled down to room temperature and 2N aq Na₂CO₃ solution is added. The mixture is stirred for 1 h and extracted several times with chloroform. The organic extracts are collected and dried (Na₂SO₄). Evaporation of the solvent and purification of the residue by reverse phase preparatory
 10 HPLC affords the desired compounds 4.

Example 163

Pyrimidinyl Compounds Having Carbon-Carbon Linkages at Position 2

- The use of diverse amidines in the cyclization reaction with suitably substituted benzoylacetates can afford pyrimidines bearing a carbon-carbon linkage in position 2.
 15 Some examples are depicted in Schemes 2 and 3 below. The desired amidines are either commercially available or can be obtained from available precursors through procedures known by one skilled in the art.



Scheme 2

6-(3-methoxyphenyl)-2-(2-morpholin-4-yl-2-oxoethyl)-3-hydropyrimidin-4-one (8)

In a round bottom flask, oven dried and kept under N₂, Cs₂CO₃ (1.5 eq) is suspended in dry DMF. 3-morpholin-4-yl-3-oxopropanamidine hydrochloride (1.2 eq) is

added, followed by ethyl ethyl 3-(3-methoxyphenyl)-3-oxopropanoate 7 (1eq). The reaction mixture is stirred at 115 °C overnight, then cooled down to room temperature. The DMF is distilled off under reduced pressure and the residue is dissolved in water, neutralizing with 5% HCl solution. The aqueous phase is then extracted with CH₂Cl₂ (x5). The organic extracts are collected and dried (Na₂SO₄). After evaporation of the solvent under reduced pressure the desired 6-(3-methoxyphenyl)-2-(2-morpholin-4-yl-2-oxoethyl)-3-hydropyrimidin-4-one 8 is obtained.

6-(3-methoxyphenyl)-2-(2-morpholin-4-yl-2-oxoethyl)pyrimidin-4-yl
(trifluoromethyl)sulfonate (9)

6-(3-methoxyphenyl)-2-(2-morpholin-4-yl-2-oxoethyl)-3-hydropyrimidin-4-one 8 (1eq) is dissolved in CH₂Cl₂ in a round bottom flask, oven dried and kept under N₂. Triethylamine is added (1.4 eq) followed by N-Phenyl trifluoromethanesulfonimide (1.2 eq) and DMAP (10 mol%). The reaction mixture is stirred at room temperature overnight. The solvent is evaporated under reduced pressure and the residue purified by chromatography on silicagel (ethyl acetate/hexanes 1:5), obtaining the desired 6-(3-methoxyphenyl)-2-(2-morpholin-4-yl-2-oxoethyl)pyrimidin-4-yl (trifluoromethyl)sulfonate 9.


2-[6-(3-methoxyphenyl)-4-(3-pyridylamino)pyrimidin-2-yl]
-1-morpholin-4-ylethan-1-one (10)

A round bottom flask, oven dried and kept under N₂ atmosphere is charged with Cs₂CO₃ (1.4 eq), Pd(OAc)₂ (5 mol%), and S-(-)-BINAP (7.5 mol %). The flask is purged with N₂ for about 5-10 min and a solution of compound 9 (1eq) in dry THF is added via a syringe, followed by 3-aminopyridine (2 eq) in one portion. The flask is equipped with a reflux condenser, purged again with N₂ for 5 min and the reaction mixture is refluxed overnight. The reaction mixture is cooled down to room temperature and the solvent is evaporated under reduced pressure. The residue is washed with water (x2) and triturated with methanol to afford the desired 2-[6-(3-methoxyphenyl)-4-(3-pyridylamino)pyrimidin-2-yl]-1-morpholin-4-ylethan-1-one 10.

[6-(3-methoxyphenyl)-2-(2-morpholin-4-ylethyl)pyrimidin-4-yl]-3-pyridylamine (11)

A dry round bottom flask is charged with LiAlH₄ (4 eq), and dry THF is added. The suspension is cooled down to 0 °C and a solution of compound 10 in THF is added

dropwise. The reaction mixture is stirred 4 h at room temperature then cooled down to 0 °C and quenched with water, followed by 10% NaOH and then water again. The mixture is stirred overnight and the solid is filtered off. The aqueous phase is extracted with CH₂Cl₂, the organic extracts are collected and dried (Na₂SO₄). After evaporation of the solvent under reduced pressure and purification by reverse phase preparatory HPLC [6-(3-methoxyphenyl)-2-(2-morpholin-4-ylethyl)pyrimidin-4-yl]-3-pyridylamine **11** is obtained.



Scheme 3

[2-(bromoethyl)-6-(3-methoxyphenyl)pyrimidin-4-yl]-3-pyridylamine (15)

[2-ethyl-6-(3-methoxyphenyl)pyrimidin-4-yl]-3-pyridylamine **14** (synthesized with a procedure similar to the one previously described) (1 eq) is dissolved in acetic acid, then sodium acetate (2 eq) is added. To this mixture, a solution of bromine (1 eq) in acetic acid is added dropwise. The reaction is stirred at room temperature for 3 h. The reaction mixture is concentrated under reduced pressure, water is added and the solution basified (pH ~10-11) with sat. aq. Na₂CO₃ solution. The product **15** crashes out, is filtered off, dried, and used as is in the following step.


[6-(3-methoxyphenyl)-2-(morpholin-4-ylethyl)pyrimidin-4-yl]-3-pyridylamine (16)

Compound **15** (1 eq) is dissolved in 3 ml of dimethyl acetamide and morpholine (5 eq) is added. The reaction mixture is stirred at 60 °C for 4 h, then cooled down to room temperature. Water is added to the mixture and the crash out is filtered off, washed with water and purified by reverse phase preparatory HPLC, obtaining [6-(3-methoxyphenyl)-2-(morpholin-4-ylethyl)pyrimidin-4-yl]-3-pyridylamine **16**.

Example 164

4-C and 4-O Substituted Pyrimidinyl Compounds

Substitution at the 4-position is not limited to an amino group, as described in Example 163. Position 4 can also bear an oxygen or a carbon linker. Ethers and 4-aryl, alkyl or 4-substituted alkyl pyrimidines can be obtained *via* standard procedures (i.e. S_NAr, Mitsunobu, Suzuki, Stille, Heck and Sonogashira couplings) known to the one skilled in the art and exemplified by the following schemes 4 and 5.



Scheme 4

10 General procedure for the synthesis of 4-Alkoxy-6-(3-methoxyphenyl)-2-morpholin-4-ylpyrimidines (18)

NaH (60% in mineral oil, ~1.2 eq) is suspended in dry NMP and the desired alcohol (1 eq) is added. The reaction mixture is stirred at room temperature for 1 h, then triflate 17 is added in one portion and the mixture heated at 100 °C for 2 h. The reaction is cooled down to room temperature, quenched with water, and heated at 60 °C. The water is extracted with CH₂Cl₂, the organic extracts are dried (Na₂SO₄) evaporated and purified by reverse phase preparatory HPLC, obtaining compound 18.

General procedure for the synthesis of 6-(3-methoxyphenyl)-2-morpholin-4-yl-4-Aryl (or heteroaryl) oxypyrimidines (20)

20 The desired hydroxy substituted aromatic or heteroaromatic compound (1 eq) and triflate 17 (1 eq) are dissolved in DMF, and solid K₂CO₃ (2 eq) is added in one portion. The reaction mixture is heated at 115 °C overnight. The reaction mixture is cooled down


to room temperature, most of the DMF is distilled off and water is added to the residue to obtain a precipitate. The solid is filtered off, dried, and purified by reverse phase preparatory HPLC, obtaining compound 19. Ester 19 is dissolved in a mixture of EtOH and 30% NaOH (1:1) and the solution is heated at 60 °C overnight. The solution is cooled
5 down to room temperature and concentrated. Purification by reverse phase preparatory HPLC affords compound 20.

General procedure for Suzuki couplings: Synthesis of 3-Methoxy-1-(2-morpholin-4-yl-6-aryl (or heteroaryl) pyrimidin-4-yl) benzenes (22)

A round bottom flask is charged with 2N Na₂CO₃ solution (4 eq) and THF and
10 the mixture is sparged with N₂ through a dispersion tube. Triflate 21 (1 eq) and the desired boronic acid or boronate (1.2 eq) are subsequently added, followed by Pd(dppf)₂Cl₂ (2.5 mol %). The reaction mixture is refluxed overnight, cooled to room temperature and diluted with EtoAc. The two phases are separated, the organic phase is washed with 2N aq Na₂CO₃, brine, and dried (Na₂SO₄). Evaporation of the solvent
15 under reduced pressure, and purification by column chromatography on silicagel affords the desired product 22.

General Procedure for Sonogashira couplings: Synthesis of 3-Methoxy-1-(2-morpholin-4-yl-6-alkynyl pyrimidin-4-yl) benzenes (23)

A round bottom flask is charged with THF, and the solvent is sparged with
20 nitrogen for 10 minutes, using a dispersion tube. The alkyne (1 eq), pyrrolidine (2 eq) and triflate 21 (1 eq) are added, while bubbling nitrogen through the solution. Pd[P(Ph)₃]₄ (2.5 mol%) is added last, and the sparging stopped. The flask is equipped with a reflux condenser and the reaction mixture is refluxed overnight under nitrogen, then cooled down to room temperature. The THF is evaporated, the residue is triturated
25 with water and ether and purified by reverse phase preparatory HPLC to obtain product 23.



Scheme 5

6-(3-methoxyphenyl)-2-morpholin-4-ylpyrimidine-4-carbonitrile (24)

A dry round bottom flask is charged with triflate 21 (1 eq) and zinc cyanide (2 eq), and DMF is added. Nitrogen is bubbled through the solution for 5 minutes and Pd[P(Ph)₃]₄ is added in one portion. The reaction mixture is stirred at 90 °C overnight. After cooling down to room temperature sat. NaHCO₃ is added, and the mixture extracted with EtOAc. The organic extracts are collected and dried (Na₂SO₄). Evaporation of the solvent under reduced pressure and purification by column chromatography on silicagel (10% methanol in methylene chloride) afforded the desired product 24.

6-(3-methoxyphenyl)-2-morpholin-4-ylpyrimidine-4-carboxylic acid (29)

Compound 24 is dissolved in a 1:1 mixture of EtOH and 30% aq. NaOH. The solution is heated to 100 °C for 2 h. The mixture is cooled down to room temperature, concentrated and neutralized with 1 N HCl. The precipitate thus formed is washed with water twice and dried, to afford the desired product 25.

Synthesis of N,N-disubstituted[6-(3-methoxyphenyl)
-2-morpholin-4-ylpyrimidin-4-yl]carboxamides (30)

Carboxylic acid 29 (1 eq) is suspended in DMF. Et₃N (2 eq) and the desired amine (1.3 eq) are added, followed by EDC (1.2 eq) and HOAT (1.2 eq). The reaction mixture is stirred at room temperature for 2 days. Water is added, and the mixture is extracted with EtOAc. The residue is purified by preparatory HPLC obtaining the desired product 30.

The above compounds can be modified further via synthetic methodologies known to the one skilled in the art. In compound 23 the triple bond can be completely or partially reduced under hydrogenation conditions by appropriate choice of the catalyst, such as Ni, 10% Pd/C, 5% Pd/C, or Lindlar catalyst. Nitrile 24 can be reduced under different conditions to the 4-aminomethyl pyrimidine 25 or to aldehyde 27, which can be functionalized further capping with carboxylic acids (on 25) or *via* reductive aminations (on both 25 and 27) to afford a variety of 4-aminomethyl substituted pyrimidines.


15

Example 165

6-Substituted Pyrimidinyl Compounds

The group in position 6 can be subject to synthetic modifications after formation of an advanced intermediate, when aromatic 1,2- 1,3- and 1,4-dialdehydes are used as substrates in the Knoevenagel condensation step, as exemplified by Scheme 6. The formyl group can then be directly reduced to hydroxymethyl, or used as a handle for reductive aminations.

20



Scheme 6

Diethyl 2-[(3-formylphenyl)methylene]propane-1,3-dioate (32)

5 Benzene 1,3 dicarbaldehyde (1 eq) is dissolved in toluene, and diethylmalonate (1 eq) is added, followed by piperidine (0.1 eq) and AcOH (0.1 eq). The flask is equipped with a Dean Stark trap and the reaction mixture is refluxed overnight. The reaction mixture is cooled down to room temperature, washed with water, 2% aq HCl, sat. aq NaHCO₃, brine, and dried. The solvent is evaporated under reduced pressure and the product 32 is isolated by column chromatography on silicagel.

10 Diethyl 2-[(3-(1,3-dioxolan-2-yl)phenyl)methylene]propane-1,3-dioate (33)

The solution of product 32 in toluene, from the previous reaction, is filtered, transferred to a round bottom flask, and ethylene glycol (2.4 eq) is added, followed by p-toulenesulfonic acid (0.5 eq). The reaction mixture is refluxed overnight with a Dean-Stark trap, then cooled down to room temperature, washed with sat. aq NaHCO₃, brine, and dried (Na₂SO₄). Product 33 is not purified further but used as is in the following step.

15

Ethyl 4-(3-(1,3-dioxolan-2-yl)phenyl)-6-morpholin-4-yl-2-oxo-
1,3,4-trihydropyridine-3-carboxylate (34)

A dry round bottom flask is charged with dry EtOH, and Na (3 eq) is added. The reaction mixture is stirred till complete dissolution of Na, then morpholino carboxamide hydrochloride (1.2 eq) is added, followed by compound 33 (1 eq). The reaction mixture is stirred at room temperature overnight. The solvent is removed under reduced pressure and water is added to the residue. The solid thus obtained is filtered, washed with water and dried, to afford crude 34 which is used in the next step without further purification.

4-(3-(1,3-Dioxolan-2-yl)phenyl)-6-morpholin-4-ylhydropyridin-2-one (36)

Substrate 34 (1 eq) is dissolved in CH₃CN. DDQ (1.2 eq) is added. The reaction mixture is stirred at room temperature overnight. The solvent is evaporated, and the residue purified by column chromatography on silicagel (10% MeOH in CH₂Cl₂) to obtain the desired ester 35, which undergoes hydrolysis and decarboxylation to 36 under conditions similar to the ones previously described for compound 20.

4-(3-(1,3-dioxolan-2-yl)phenyl)-6-morpholin-4-yl
-2-pyridyl (trifluoromethyl)sulfonate (37)

The title compound is prepared following the same procedure as the one used for compound 9, described above.

[4-(3-(1,3-Dioxolan-2-yl)phenyl)-6-morpholin-4-yl(2-pyridyl)]-3-pyridylamine (38)

A round bottom flask, oven dried and kept under N₂ atmosphere is charged with Cs₂CO₃ (1.4 eq), Pd(OAc)₂ (5 mol%), and S-(-)-BINAP (7.5 mol %). The flask is purged with N₂ for about 5-10 min and a solution of compound 37 (1eq) in dry THF is added via a syringe, followed by 3-aminopyridine (2 eq) in one portion. The flask is equipped with a reflux condenser, purged again with N₂ for 5 min and the reaction mixture is refluxed overnight. The reaction mixture is cooled down to room temperature and the solvent is evaporated under reduced pressure. The residue is washed with water (x2) and triturated with methanol to afford compound 38.

3-[2-morpholin-4-yl-6-(3-pyridylamino)-4-pyridyl]benzaldehyde (39)


Compound 38 was suspended in wet acetone and p-toluenesulfonic acid (0.2 eq) was added. The reaction mixture was refluxed overnight, then concentrated under

reduced pressure. The residue was triturated with diethyl ether, water and MeOH to afford the desired aldehyde **39**.

{3-[2-morpholin-4-yl-6-(3-pyridylamino)-4-pyridyl]phenyl}methan-1-ol (**40**)

Aldehyde **39** is suspended in THF and DIBALH (1.6 N solution in THF, 3 eq) is added dropwise via a syringe. The reaction mixture is stirred at room temperature overnight, then quenched with water. The aqueous phase is extracted repeatedly with EtOAc. The organic extracts are collected and dried (Na₂SO₄). Evaporation of the solvent under reduced pressure and purification by reverse phase preparatory HPLC, affords compound **40**.

Substituents in position 6 can be varied using suitable aldehydes in the Knoevenagel step, as previously described. These substrates need not be limited to aromatic aldehydes. The synthetic route can be extended to heteroaromatic, heterocyclic and aliphatic aldehydes. In particular, substituted phenylacetaldehydes (**41**, n=1) and 3-phenylpropionaldehydes (**41**, n=2), allow access to compounds such as **42** in which the substituted aromatic group is linked to the pyrimidine core by a single carbon or a 2 carbons spacer, as shown in Scheme 7.



Scheme 7

Example 166

PI3K Assay Procedures

Method 1: Homogenous solution phase assay

Compounds to be tested are dissolved in DMSO and directly distributed to 384-well flashplates at 1.25 μ L per well. To start the reaction, 20 μ L of 6 nM PI3 kinase are added into each well followed by 20 μ L of 400 nM ATP containing a trace of radio-labeled ATP and 900 nM 1-alpha-phosphatidylinositol (PI). The plates are briefly centrifuged to remove any air gap. The reaction is performed for 15 minutes and then stopped by the addition of 20 μ L of 100 mM EDTA. The stopped reaction is incubated

overnight at RT to allow the lipid substrate to bind by hydrophobic interaction to the surface of the flashplate. The liquid in the wells is then washed away, and the labeled substrate is detected with scintillation counting.

Method 2: One step solid phase assay

5 This method is similar to Method 1 except that the lipid substrate (1-alpha-phosphatidylinositol) is first dissolved in a coating buffer and incubated on flashplate at room temperature over night to allow the lipid substrate to bind by hydrophobic interaction to the surface of the flashplate. Unbound substrate is then washed away. On the day of assay, 20 μ L of 6 nM PI3 kinase are added into each well followed by 20 μ L of
10 400 nM ATP containing trace of radio-labeled ATP. Compounds are added together with enzyme and ATP to the lipid-coated plates. The plates are briefly centrifuged to remove any air gap. The reaction is performed for two to three hours. The reaction is stopped by addition of 20 μ L of 100 mM EDTA or by immediate plate washing. Phosphorylated lipid substrate is detected with scintillation counting.

15 The compounds of Examples 11a, 13, 19, 34-49, 51-53, 55, 57-59, 61-64, 68, 71-76, 79, 81, 82, 85-87, 89-91, 118, 119, 121, 122, 124 and 133-156 displayed an IC₅₀ value of less than 20 μ M with respect to PI3K when tested in the homogeneous solution assay (Method 1), as described above. The compounds of Examples 20, 21, 23, 47,
20 55-60, 62, 63, 65, 70, 71-75, 77-95, 97-120, 122-125, 127, 129, 130, 133, 137 and 143-155 displayed an IC₅₀ value of less than 20 μ M with respect to PI3K when tested in the one step solid phase assay (Method 2), as described above.


 It should be understood that the organic compounds according to the invention may exhibit the phenomenon of tautomerism. As the chemical structures within this specification can only represent one of the possible tautomeric forms, it should be
25 understood that the invention encompasses any tautomeric form of the drawn structure.

 It is understood that the invention is not limited to the embodiments set forth herein for illustration, but embraces all such forms thereof as come within the scope of the above disclosure.

 While the preferred embodiment of the invention has been illustrated and
30 described, it will be appreciated that various changes can be made therein without departing from the spirit and scope of the invention.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A compound having the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,

- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and

R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^z is H or substituted or unsubstituted alkyl group; and

R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C₁-C₆-alkyl; and r is 0, 1, or 2;

with the proviso that when X is O, then Y is substituted or unsubstituted aryl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heteroaryl;

with the proviso that when W is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido morpholino, piperazino, or N-substituted piperazino, R₂ is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido-thiomorpholino, piperazino, or N'-[acetyl(alkanoyl of 1 to 3 carbon atoms)]piperazino, and X is NH, then Y is not hydrogen, alkyl of 1 to 3 carbon atoms, cyclohexyl, phenyl, chloro-phenyl, carboxy-phenyl, carbomethoxy-phenyl, or pyridyl;

with the proviso that when W is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido morpholino, piperazino, or N-substituted piperazino, R₂ is morpholino, thiomorpholino, 1-oxido-thiomorpholino, 1,1-dioxido-thiomorpholino, piperazino, or N'-[acetyl(alkanoyl of 1 to 3 carbon atoms)]piperazino, and X is a direct link, then Y is not phenyl, substituted or unsubstituted C₁-C₆-alkyl, or 1-oxidothiomorpholino; and

with the proviso that when R₂ is phenyl independently substituted with one to five substituents selected from hydrogen, cycloalkyl, heterocycloalkyl, halo, nitro, amino, sulphonamido, or alkylsulphonylamino, R₁ is hydrogen, haloalkyl, alkyl, or halo, and X is


NR^{1x} , then Y is substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocyclyl.

2. The compound of claim 1, wherein

Y is selected from the group consisting of


- (1) substituted or unsubstituted $\text{C}_1\text{-C}_6$ -alkyl,
- (2) substituted or unsubstituted aryl,
- (3) substituted or unsubstituted heterocyclyl, and
- (4) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) $-\text{N}(\text{R}^{1x})-$,
- (3) $-(\text{CH}_2)_m-\text{C}(\text{R}^{2x}, \text{R}^{3x})-\text{N}(\text{R}^{1x})-$, and
- (4) ,

wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted $\text{C}_1\text{-C}_6$ -alkyl; and

W is selected from the group consisting of



wherein Z is $-\text{O}-$ or $-\text{NR}^z-$, wherein R^{4w} is H or substituted or unsubstituted $\text{C}_1\text{-C}_6$ -alkyl.

3. The compound of claim 1, wherein


Y is selected from the group consisting of

- (1) substituted or unsubstituted heterocyclyl,
- (2) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of


- (1) a direct link,
- (2) $-\text{N}(\text{R}^{1x})-$,

(3) $-(\text{CH}_2)_m-\text{C}(\text{R}^{2x}, \text{R}^{3x})-\text{N}(\text{R}^{1x})-$, and



wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl; and

W is selected from the group consisting of



wherein Z is $-\text{O}-$ or $-\text{NR}^z-$, wherein R^{4w} is H or substituted or unsubstituted C_1 - C_6 -alkyl.

4. The compound of claim 1, wherein


Y is substituted or unsubstituted aryl;

X is selected from the group consisting of

(1) a direct link,


(2) $-\text{N}(\text{R}^{1x})-$,

(3) $-(\text{CH}_2)_m-\text{C}(\text{R}^{2x}, \text{R}^{3x})-\text{N}(\text{R}^{1x})-$, and



wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl; and

W is selected from the group consisting of




wherein Z is $-\text{O}-$ or $-\text{NR}^z-$, wherein R^{4w} is H or substituted or unsubstituted C_1 - C_6 -alkyl.

5. The compound of claim 1, wherein


Y is substituted or unsubstituted alkyl;

X is selected from the group consisting of

- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and
- (4) ,

wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl; and

W is selected from the group consisting of




wherein Z is $-O-$ or $-NR^Z-$, wherein R^{4w} is H or substituted or unsubstituted C_1 - C_6 -alkyl.

6. The compound of claim 1, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted heterocyclyl,
- (2) substituted or unsubstituted heteroaryl;


X is selected from the group consisting of

- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and
- (4) ,

wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl;

R_2 is substituted or unsubstituted aryl; and

W is




wherein Z is -O- or -NH-.

7. The compound of claim 1, wherein

Y is substituted or unsubstituted aryl;

X is selected from the group consisting of


- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and



wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl;

R_2 is substituted or unsubstituted aryl; and

W is




wherein Z is -O- or -NH-.

8. The compound of claim 1, wherein

Y is substituted or unsubstituted alkyl;

X is selected from the group consisting of


- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and



wherein R^{1x} , R^{2x} , R^{3x} are independently H or substituted or unsubstituted C_1 - C_6 -alkyl;


R_2 is substituted or unsubstituted aryl; and

W is



wherein Z is -O- or -NH-.

9. The compound of claim 1, having the formula II:



(II)


wherein Y is selected from the group consisting of

- (1) substituted or unsubstituted C_1 - C_6 -alkyl,
- (2) substituted or unsubstituted aryl,
- (3) substituted or unsubstituted heterocyclyl, and
- (4) substituted or unsubstituted heteroaryl; and


X is selected from the group consisting of

- (1) a direct link,
- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$, and
- (4) .


10. The compound of claim 1, having the formula II:




wherein Y and X, taken together, are selected from the group consisting of




11. The compound of claim 1, having the formula II:



wherein Y and X, taken together, are selected from the group consisting of




12. The compound of claim 1, having the formula II:




(II)

wherein, Y and X, taken together, are selected from the group consisting of



13. The compound of claim 1, having the formula III:




(III)

wherein R_3 , R_4 , R_5 , R_6 are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C_1 - C_6 -alkyl,
- (3) $-COOR^{1t}$,
- (4) $-COONH_2$,
- (5) $-OR^{1t}$, and
- (6) $-NHR^{1t}$.

14. The compound of claim 1, having the formula IV:




(IV)

wherein R₃, R₄, R₅, R₆ are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},
- (4) -COONH₂,
- (5) -OR^{1t}, and
- (6) -NHR^{1t}.

15. The compound of claim 1, having the formula V:



(V)

wherein R₃, R₄, R₅, R₆ are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},
- (4) -COONH₂,
- (5) -OR^{1t}, and
- (6) -NHR^{1t}; and

R^{2a} and R^{2b} are selected from the group consisting of


- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) halo,
- (4) $-(CH_2)_q-N(R^{2c}, R^{2d})$,
- (5) $-(CH_2)_q-N(R^{2c}, R^{2d})COR^{2e}$,
- (6) $-(CH_2)_q-OR^{2e}$,
- (7) $-(CH_2)_q-OCOR^{2e}$,
- (8) $-(CH_2)_q-OCOOR^{2e}$,
- (9) $-(CH_2)_q-COOR^{2e}$,
- (10) $-(CH_2)_q-CONR^{2c}$,
- (11) $-CN$,
- (12) $-NO_2$,
- (13) $-SO_2NH_2$,
- (14) $-NHSO_2CH_3$, and
- (15) $-SO_2R^{2f}$,

wherein R^{2c} , R^{2d} , R^{2e} , and R^{2f} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted phenyl; and


q is 0, 1, 2, 3, or 4.

16. The compound of claim 1, having the formula VI:




(VI)

wherein R_2 is selected from the group consisting of



17. The compound of claim 1, having the formula VII:




(VII)

wherein R₇, R₈, R₉, and R₁₀ are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},
- (4) -COONH₂,
- (5) -OR^{1t}, and
- (6) -NHR^{1t}.

18. The compound of claim 1, having the formula VIII:




(VIII)

wherein R₇, R₈, R₉, R₁₀ are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},
- (4) -CONH₂,
- (5) -OR^{1t}, and
- (6) -NHR^{1t}.

19. The compound of claim 1, having the formula IX:



(IX)

wherein R^{1a} and R^{1b} are selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) halo,
- (4) $-(CH_2)_q-N(R^{2c}, R^{2d})$,
- (5) $-(CH_2)_q-N(R^{2c}, R^{2d})COR^{2e}$,
- (6) $-(CH_2)_q-OR^{2e}$,
- (7) $-(CH_2)_q-OCOR^{2e}$,
- (8) $-(CH_2)_q-OCOOR^{2e}$,
- (9) $-(CH_2)_q-COOR^{2e}$,
- (10) $-(CH_2)_q-CONR^{2c}$,
- (11) -CN,
- (12) -NO₂,
- (13) -SO₂NH₂,
- (14) -NHSO₂CH₃, and
- (15) -SO₂R^{2f},

wherein R^{2c} , R^{2d} , R^{2e} , and R^{2f} are selected from the group consisting of


- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted phenyl; and

wherein R_7 is selected from the group consisting of


- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOR^{1t},

- (4) $-\text{CONH}_2$,
- (5) $-\text{OR}^{1t}$, and
- (6) $-\text{NHR}^{1t}$.


20. The compound of claim 1, having the formula X:



wherein R_2 is selected from the group consisting of





21. The compound of claim 1, having the formula XI:



wherein R^{2g} is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) $-\text{CONHR}^{2h}$,
- (4) $-\text{CON}(R^{2h})-(\text{CH}_2)_{2-3}-\text{N}(R^{2h}, R^{2i})$,
- (5) $-\text{COR}^{2j}$,
- (6) $-\text{CO}_2R^{2j}$,
- (7) $-\text{COC}_1\text{-C}_6\text{-alkyl-CO}_2\text{H}$,
- (8) $-\text{CH}_2\text{-OC}(=\text{O})R^{2i}$,
- (9) $-\text{CH}_2\text{-OC}(=\text{O})\text{NHCHR}^{2i}\text{CO}_2R^{2j}$,
- (10) $-\text{P}(=\text{O})(\text{OR}^{2k}, \text{OR}^{2p})$,




wherein R^{2h} , R^{2i} , R^{2j} , R^{2k} , and R^{2p} are selected from the group consisting

of

- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted aryl.



22. The compound of claim 1, having the formula XII:



(XII)

wherein R^{2g} is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted alkyl,
- (3) -CONHR^{2h},
- (4) -CON(R^{2h})-(CH₂)₂₋₃-N(R^{2h}, R²ⁱ),
- (5) -COR^{2j},
- (6) -CO₂R^{2j},
- (7) -COC₁-C₆-alkyl-CO₂H,
- (8) -CH₂-OC(=O)R²ⁱ,
- (9) -CH₂-OC(=O)NHCHR²ⁱCO₂R^{2j},
- (10) -P(=O)(OR^{2k}, OR^{2p}),




wherein R^{2h}, R²ⁱ, R^{2j}, R^{2k}, and R^{2p} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted alkyl, and
- (c) substituted or unsubstituted aryl.

23. A composition, comprising a pharmaceutically acceptable carrier and an amount of a compound effective to inhibit phosphatidylinositol (PI) 3-kinase activity in a

human or animal subject when administered thereto, wherein the compound has the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,

- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and

R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,
- (c) -S-,

- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^z is H or substituted or unsubstituted alkyl group; and
R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},


wherein R^{5w} is H or C₁-C₆-alkyl; and

r is 0, 1, or 2.

24. The composition of Claim 23 further comprising at least one additional agent for the treatment of cancer.

25. The composition of Claim 24, wherein the at least one additional agent for the treatment of cancer is selected from irinotecan, topotecan, gemcitabine, gleevec, herceptin, 5-fluorouracil, leucovorin, carboplatin, cisplatin, taxanes, tezacitabine, cyclophosphamide, vinca alkaloids, imatinib, anthracyclines, rituximab, tamoxifen, CPT 11, and trastuzumab.

26. A method for treating a condition by modulation of phosphatidylinositol (PI) 3-kinase activity comprising administering to a human or animal subject in need of such treatment an effective amount of a compound having the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and

(6) $-\text{NHR}^{1t}$,


wherein R^{1t} is H or $\text{C}_1\text{-C}_6\text{-alkyl}$;

R_2 is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted $\text{C}_1\text{-C}_6\text{-alkyl}$,
- (2) $-\text{N}(\text{R}^{1w}, \text{R}^{2w})$, and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted $\text{C}_1\text{-C}_6\text{-alkyl}$,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and

R^{2w} are not both H;

Z is selected from the group consisting of

- (a) $-\text{O}-$,
- (b) $-\text{NR}^z$,
- (c) $-\text{S}-$,
- (d) $-\text{SO}-$,
- (e) $-\text{SO}_2-$, and
- (f) $-\text{CH}_2-$,

wherein R^z is H or substituted or unsubstituted alkyl group; and

R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted $\text{C}_1\text{-C}_6\text{-alkyl}$,
- (c) $-\text{COOR}^{5w}$,
- (d) $-\text{CONH}_2$,
- (e) $-\text{OR}^{5w}$, and

(f) $-\text{NHR}^{5w}$,


wherein R^{5w} is H or $\text{C}_1\text{-C}_6\text{-alkyl}$; and

r is 0, 1, or 2.

27. The method of Claim 26, wherein the compound has an IC_{50} value of less than about $20\ \mu\text{M}$ in a cell proliferation assay.

28. The method of Claim 26, wherein the condition is cancer.

29. A method for inhibiting phosphatidylinositol (PI) 3-kinase activity in a human or animal subject, comprising administering to the human or animal subject a composition comprising an amount of a compound effective to inhibit phosphatidylinositol (PI) 3-kinase activity in the human or animal subject, wherein the compound has the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted $\text{C}_1\text{-C}_6\text{-alkyl}$,
- (2) substituted or unsubstituted $\text{C}_2\text{-C}_6\text{-alkenyl}$,
- (3) substituted or unsubstituted $\text{C}_2\text{-C}_6\text{-alkynyl}$,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) $-\text{N}(\text{R}^{1x})-$,
- (3) $-(\text{CH}_2)_m-\text{C}(\text{R}^{2x}, \text{R}^{3x})-\text{N}(\text{R}^{1x})-$,
- (4) $-\text{O}-$,

- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^Z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^Z is H or substituted or unsubstituted alkyl group; and


R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C_1 - C_6 -alkyl; and

r is 0, 1, or 2.

30. A method for treating a cancer disorder in a human or animal subject, comprising administering to the human or animal subject a composition comprising an amount of a compound effective to inhibit phosphatidylinositol (PI) 3-kinase activity in the human or animal subject, wherein the compound has the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^Z is H or substituted or unsubstituted alkyl group; and R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) $-COOR^{5w}$,
- (d) $-CONH_2$,
- (e) $-OR^{5w}$, and
- (f) $-NHR^{5w}$,


wherein R^{5w} is H or C_1 - C_6 -alkyl; and

r is 0, 1, or 2.

31. The method of Claim 30 further comprising administering to the human or animal subject at least one additional agent for the treatment of cancer.

32. The method of Claim 31, wherein the at least one additional agent for the treatment of cancer is selected from irinotecan, topotecan, gemcitabine, gleevec, herceptin, 5-fluorouracil, leucovorin, carboplatin, cisplatin, taxanes, tezacitabine, cyclophosphamide, vinca alkaloids, imatinib, anthracyclines, rituximab, tamoxifen, CPT 11, and trastuzumab.

33. A method for inhibiting tumor growth in a human or animal subject, comprising administering to the human or animal subject in need thereof an effective amount of a compound having the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C_1 - C_6 -alkyl,
- (2) substituted or unsubstituted C_2 - C_6 -alkenyl,

- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and

R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^z is H or substituted or unsubstituted alkyl group; and


R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C₁-C₆-alkyl; and

r is 0, 1, or 2.

34. A method for inhibiting the proliferation of capillaries in a human or animal subject, comprising administering to the human or animal subject in need thereof an effective amount of a compound having the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,

- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,

- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^z is H or substituted or unsubstituted alkyl group; and


R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C₁-C₆-alkyl; and

r is 0, 1, or 2.

35. A compound for use in the treatment of cancer, wherein the compound has the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,

- (2) $-N(R^{1x})-$,
- (3) $-(CH_2)_m-C(R^{2x}, R^{3x})-N(R^{1x})-$,
- (4) $-O-$,
- (5) $-S-$,
- (6) $-SO-$,
- (7) $-SO_2-$,
- (8) $-C(R^{2x}, R^{3x})-$, and
- (9) ,

wherein R^{1x} , R^{2x} , and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1-C_6 -alkyl,
- (c) substituted or unsubstituted C_2-C_6 -alkenyl,
- (d) substituted or unsubstituted C_2-C_6 -alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R_1 is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C_1-C_6 -alkyl,
- (3) $-COOH$,
- (4) halo,
- (5) $-OR^{1t}$, and
- (6) $-NHR^{1t}$,


wherein R^{1t} is H or C_1-C_6 -alkyl;

R_2 is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C_1-C_6 -alkyl,
- (2) $-N(R^{1w}, R^{2w})$, and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^Z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^Z is H or substituted or unsubstituted alkyl group; and


R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C_1 - C_6 -alkyl; and

r is 0, 1, or 2.

34. A method for inhibiting the proliferation of capillaries in a human or animal subject, comprising administering to the human or animal subject in need thereof an effective amount of a compound having the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) substituted or unsubstituted C₂-C₆-alkenyl,
- (3) substituted or unsubstituted C₂-C₆-alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) -N(R^{1x})-,
- (3) -(CH₂)_m-C(R^{2x}, R^{3x})-N(R^{1x})-,
- (4) -O-,
- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) -NR^z-,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,


wherein R^z is H or substituted or unsubstituted alkyl group; and R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) $-\text{COOR}^{5w}$,
- (d) $-\text{CONH}_2$,
- (e) $-\text{OR}^{5w}$, and
- (f) $-\text{NHR}^{5w}$,

wherein R^{5w} is H or C_1 - C_6 -alkyl; and

r is 0, 1, or 2.

36. Use of a compound in the manufacture of a medicament for the treatment of cancer, wherein the compound has the formula I:




or a stereoisomer, tautomer, pharmaceutically acceptable salt, ester, or prodrug thereof, wherein

Y is selected from the group consisting of

- (1) substituted or unsubstituted C_1 - C_6 -alkyl,
- (2) substituted or unsubstituted C_2 - C_6 -alkenyl,
- (3) substituted or unsubstituted C_2 - C_6 -alkynyl,
- (4) substituted or unsubstituted aryl,
- (5) substituted or unsubstituted heterocyclyl, and
- (6) substituted or unsubstituted heteroaryl;

X is selected from the group consisting of

- (1) a direct link,
- (2) $-\text{N}(\text{R}^{1x})-$,
- (3) $-(\text{CH}_2)_m-\text{C}(\text{R}^{2x}, \text{R}^{3x})-\text{N}(\text{R}^{1x})-$,
- (4) $-\text{O}-$,

- (5) -S-,
- (6) -SO-,
- (7) -SO₂-,
- (8) -C(R^{2x}, R^{3x})-, and
- (9) ,

wherein R^{1x}, R^{2x}, and R^{3x} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C₁-C₆-alkyl,
- (c) substituted or unsubstituted C₂-C₆-alkenyl,
- (d) substituted or unsubstituted C₂-C₆-alkynyl,
- (e) substituted or unsubstituted aryl,
- (f) substituted or unsubstituted heterocyclyl,
- (g) substituted or unsubstituted heteroaryl; and

m is 0, 1, 2, 3, or 4;

R₁ is selected from the group consisting of

- (1) H,
- (2) substituted or unsubstituted C₁-C₆-alkyl,
- (3) -COOH,
- (4) halo,
- (5) -OR^{1t}, and
- (6) -NHR^{1t},


wherein R^{1t} is H or C₁-C₆-alkyl;

R₂ is selected from the group consisting of

- (1) substituted or unsubstituted aryl,
- (2) substituted or unsubstituted heteroaryl, and
- (3) substituted or unsubstituted heterocyclyl; and

W is selected from the group consisting of

- (1) substituted or unsubstituted C₁-C₆-alkyl,
- (2) -N(R^{1w}, R^{2w}), and



wherein R^{1w} and R^{2w} are selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) substituted or unsubstituted aryl,
- (d) substituted or unsubstituted heterocyclyl, and
- (e) substituted or unsubstituted heteroaryl, wherein R^{1w} and R^{2w} are not both H;

Z is selected from the group consisting of

- (a) -O-,
- (b) - NR^Z -,
- (c) -S-,
- (d) -SO-,
- (e) -SO₂-, and
- (f) -CH₂-,

wherein R^Z is H or substituted or unsubstituted alkyl group; and

R^{4w} is selected from the group consisting of

- (a) H,
- (b) substituted or unsubstituted C_1 - C_6 -alkyl,
- (c) -COOR^{5w},
- (d) -CONH₂,
- (e) -OR^{5w}, and
- (f) -NHR^{5w},

wherein R^{5w} is H or C_1 - C_6 -alkyl; and

r is 0, 1, or 2.

INTERNATIONAL SEARCH REPORT

Int: ial Application No
PCT/US 03/37294

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D407/14 C07D403/12 C07D407/12 C07D239/48
 C07D417/12 C07D403/04 C07D403/14 C07D401/14 A61K31/506
 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BUNDY G L ET AL: "SYNTHESIS OF NOVEL 2,4-DIAMINOPYRROLO-2,3-DIPYRIMIDINES WITH ANTIOXIDANT, NEUROPROTECTIVE, AN ANTIASTHMA ACTIVITY" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 38, no. 21, 1995, pages 4161-4163, XP001106595 ISSN: 0022-2623 scheme 1, compound 1 --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *G* document member of the same patent family

Date of the actual completion of the international search

1 April 2004

Date of mailing of the international search report

16/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Schuemacher, A

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/US 03/37294

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J.E. CABAJ, P.G.M. WUTS, K.E. HENEGAR: "Bromine-mediated addition of nucleophiles to the electron-rich Pyrimidine subunit of Tirilazad" J.ORG.CHEM., vol. 59, 1994, pages 5090-5092, XP002275310 compounds 1a and 1c scheme 1, compounds 3,7,8,2a and 2g ---	1
X	G.L.BUNDY, T.M.SCHWARTZ, J.R.PALMER, L.S. BANITT AND W. WATT: "Synthesis of 2,4-diaminopyrrolo[2,3-d]pyrimidines " J.HETEROCYCLIC CHEM., vol. 37, 2000, pages 1471-1477, XP002275311 scheme 1, compound 2 scheme 2, compounds 13,14 and 15 ---	1
X	D. BROWN AND G.A.R. KON: "some heterocyclic analogues of stilbenes" J.CHEM.SOC., 1948, pages 2147-2153, XP009028506 2,6-diethylpyridino-4-methylpyrimidine p.2151 ---	1
X	A.A. ABOU OUF, M.M. EL-KERDAWY: "preparation of some methyl pyrimidines" EGYPT. J. PHARM.SCI., vol. 14, no. 2, 1973, pages 180-195, XP009028502 example VA; table 2 ---	1,2,5,8, 9
X	KOWALENSKI A, STREKOWSKI L.: "unfused heterobicycles as amplifiers of Phleomycin" AUSTRA. J CHEM., vol. 34, no. 12, 1981, pages 2929-2933, XP009028501 examples 3(a)-3(c) page 2630 ---	1
X	WO 01 072745 A (CYCLACEL LIMITED, UK) 4 October 2001 (2001-10-04) page 15, line 27 - line 29; claims 1,17-20,23-25; examples 57-70 ---	1,33-37
X	WO 02 022608 A (VERTEX PHARMACEUTICALS INCORPORATED, USA) 21 March 2002 (2002-03-21) claims 1,8-10,25; table 6 ---	1,33-37
X	WO 02 062789 A (VERTEX PHARMACEUTICALS INCORPORATED, USA) 15 August 2002 (2002-08-15) claims 1,15-17; table 7 ---	1,33-37

-/-

INTERNATIONAL SEARCH REPORT

Int onal Application No

PCT/US 03/37294

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01 83456 A (IMP CANCER RES TECHNOLOGY LTD ;HAYAKAWA MASAHIKO (JP); MORITOMO HI) 8 November 2001 (2001-11-08)	1-37
Y	-& EP 1 277 738 A (YAMANOUCHI PHARMACEUTICAL CO LTD.) 22 January 2003 (2003-01-22) the whole document	1-37
P,X	WO 03 030909 A (BAYER CORPORATION, USA) 17 April 2003 (2003-04-17) claims 9,10,12-14; examples 314,315; table 5	1,33-37
P,X	WO 02 102313 A (BRISTOL-MYERS SQUIBB COMPANY, USA) 27 December 2002 (2002-12-27) claim 1; tables A,L,M	1,33-37

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 03/37294

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 26-34 and 36 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 1-9 and 23-37 relate inter alia to "pharmaceutically acceptable ester" and "prodrug", i.e. to compounds defined by reference to a desirable characteristic or property, whereas the application appears to provide no support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for said pharmaceutically acceptable ester and prodrug (see p.36, 1.16-31).

Additionally, the claims also lack clarity (Article 6 PCT) due to the use of unclear expression like "unsubstituted or substituted" which implies that the claims relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed (e.g. in all the examples W is always morpholino).

Furthermore, claim 27 does not meet the requirements of Article 6 PCT in that the matter for which protection is sought is not clearly defined: this claim attempts to define the subject-matter in terms of a specific parameter, i.e. the IC50 value.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely those parts relating to the compounds defined by formula I with the exception of any pharmaceutically acceptable ester and any prodrugs thereof as well as the compounds prepared in the examples.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 03/37294

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 01072745	A	04-10-2001	AU 4262901 A 08-10-2001
			CA 2401748 A1 04-10-2001
			CN 1420884 T 28-05-2003
			EP 1274705 A1 15-01-2003
			WO 0172745 A1 04-10-2001
			GB 2361236 A , B 17-10-2001
			HU 0300382 A2 28-06-2003
			JP 2003528872 T 30-09-2003
			US 2003149057 A1 07-08-2003
			US 2002019404 A1 14-02-2002
WO 02022608	A	21-03-2002	AU 9091201 A 26-03-2002
			AU 9091401 A 26-03-2002
			AU 9094401 A 26-03-2002
			AU 9101301 A 26-03-2002
			AU 9267001 A 26-03-2002
			AU 9455801 A 26-03-2002
			AU 9687101 A 26-03-2002
			AU 9687501 A 26-03-2002
			BR 0114088 A 17-06-2003
			CA 2422299 A1 21-03-2002
			CA 2422354 A1 21-03-2002
			CA 2422367 A1 21-03-2002
			CA 2422371 A1 21-03-2002
			CA 2422377 A1 21-03-2002
			CA 2422378 A1 21-03-2002
			CA 2422379 A1 21-03-2002
			CA 2422380 A1 21-03-2002
			CN 1469874 T 21-01-2004
			CN 1469875 T 21-01-2004
			CN 1473161 T 04-02-2004
			EP 1317447 A1 11-06-2003
			EP 1317444 A1 11-06-2003
			EP 1317448 A1 11-06-2003
			EP 1318997 A1 18-06-2003
			EP 1317449 A1 11-06-2003
			EP 1317450 A1 11-06-2003
			EP 1317452 A1 11-06-2003
			EP 1318814 A2 18-06-2003
			HU 0302172 A2 29-09-2003
			HU 0302173 A2 29-09-2003
			HU 0302411 A2 28-11-2003
			NO 20031188 A 13-05-2003
			NO 20031189 A 13-05-2003
			NO 20031190 A 13-05-2003
			NO 20031191 A 13-05-2003
			WO 0222603 A1 21-03-2002
			WO 0222601 A1 21-03-2002
			WO 0222604 A1 21-03-2002
			WO 0222605 A1 21-03-2002
			WO 0222606 A1 21-03-2002
			WO 0222607 A1 21-03-2002
			WO 0222608 A1 21-03-2002
			WO 0222602 A2 21-03-2002
			US 2003073687 A1 17-04-2003
			US 2003083327 A1 01-05-2003
			US 2003064981 A1 03-04-2003
			US 2003064982 A1 03-04-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 03/37294

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 02022608	A	US 2003055044 A1 US 2003078166 A1	20-03-2003 24-04-2003
WO 02062789	A	15-08-2002	
		AU 3116602 A	01-07-2002
		AU 3404702 A	01-07-2002
		AU 9091201 A	26-03-2002
		AU 9091401 A	26-03-2002
		AU 9094401 A	26-03-2002
		AU 9101301 A	26-03-2002
		AU 9267001 A	26-03-2002
		AU 9455801 A	26-03-2002
		AU 9687101 A	26-03-2002
		AU 9687501 A	26-03-2002
		BR 0114088 A	17-06-2003
		BR 0116411 A	11-11-2003
		BR 0116493 A	30-09-2003
		CA 2422299 A1	21-03-2002
		CA 2422354 A1	21-03-2002
		CA 2422367 A1	21-03-2002
		CA 2422371 A1	21-03-2002
		CA 2422377 A1	21-03-2002
		CA 2422378 A1	21-03-2002
		CA 2422379 A1	21-03-2002
		CA 2422380 A1	21-03-2002
		CA 2432129 A1	25-07-2002
		CA 2432131 A1	01-08-2002
		CA 2432132 A1	01-08-2002
		CA 2432222 A1	15-08-2002
		CA 2432223 A1	06-09-2002
		CA 2432303 A1	29-08-2002
		CA 2432799 A1	27-06-2002
		CA 2432872 A1	27-06-2002
		CN 1469874 T	21-01-2004
		CN 1469875 T	21-01-2004
		CN 1473161 T	04-02-2004
		EP 1345922 A1	24-09-2003
		EP 1355905 A1	29-10-2003
		EP 1317447 A1	11-06-2003
		EP 1317444 A1	11-06-2003
		EP 1317448 A1	11-06-2003
		EP 1318997 A1	18-06-2003
		EP 1317449 A1	11-06-2003
		EP 1317450 A1	11-06-2003
		EP 1317452 A1	11-06-2003
		EP 1318814 A2	18-06-2003
		EP 1345925 A2	24-09-2003
		EP 1345928 A2	24-09-2003
		EP 1345926 A2	24-09-2003
		EP 1353916 A2	22-10-2003
		EP 1345929 A2	24-09-2003
		EP 1345927 A1	24-09-2003
		HU 0302172 A2	29-09-2003
WO 0183456	A	08-11-2001	
		AU 5261001 A	12-11-2001
		CA 2407593 A1	08-11-2001
		CN 1426398 T	25-06-2003
		EP 1277738 A1	22-01-2003
		WO 0183456 A1	08-11-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 03/37294

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03030909	A	17-04-2003	WO 03030909 A1	17-04-2003
WO 02102313	A	27-12-2002	CA 2443835 A1	07-11-2002
			CA 2444436 A1	07-11-2002
			CA 2450724 A1	27-12-2002
			CA 2450934 A1	27-12-2002
			CA 2450936 A1	27-12-2002
			EP 1383743 A2	28-01-2004
			EP 1383506 A2	28-01-2004
			EP 1397142 A2	17-03-2004
			WO 02088079 A2	07-11-2002
			WO 02088080 A2	07-11-2002
			WO 02087513 A2	07-11-2002
			WO 02102313 A2	27-12-2002
			WO 02102314 A2	27-12-2002
			WO 02102315 A2	27-12-2002
			US 2003104974 A1	05-06-2003
			US 2003092908 A1	15-05-2003
			US 2003092721 A1	15-05-2003
			US 2003162802 A1	28-08-2003
			US 2003100571 A1	29-05-2003
			US 2003191143 A1	09-10-2003